



Introduction to Lebesgue Measure

Prelude

There were two problems with calculus: there are functions where

- $f(x) \neq \int f'(x) dx$ (c.f., O-G #30)
- $f(x) \neq \frac{d}{dx} \left[\int f(x) dx \right]$

In his 1902 dissertation, “*Intégrale, long-ueur, aire,*” Lebesgue wrote, “It thus seems to be natural to search for a definition of the integral which makes integration the inverse operation of differentiation in as large a range as possible.”



Henri Lebesgue's Mathematical Genealogy (partial)

What's in a Measure

Goals

THE BEST *measure* would be a real-valued set function μ that satisfies

- 1 $\mu(I) = \text{length}(I)$ where I is an interval
- 2 μ is *translation invariant*: $\mu(x + E) = \mu(E)$ for any $x \in \mathbb{R}$
- 3 if $\{E_n\}$ is pairwise disjoint, then $\mu(\bigcup_n E_n) = \sum_n \mu(E_n)$
- 4 $\text{dom}(\mu) = \mathcal{P}(\mathbb{R})$ (the power set of \mathbb{R})

THE BAD NEWS:

$$\left\{ \begin{array}{l} \text{continuum hypothesis} \\ + \text{axiom choice} \end{array} \right\} \implies 1, 3, \text{ and } 4 \text{ are incompatible}$$

THE PLAN:

- Give up on 4. (cf. *Vitali*)
- 1. and 2. are nonnegotiable
- Weaken 3., then reclaim it

Sigma Algebras

Definition

Sigma Algebra of Sets

Algebra: A collection of sets \mathcal{A} is an *algebra* iff \mathcal{A} is closed under unions and complements.

σ -Algebra: An algebra of sets \mathcal{A} is a σ -*algebra* iff \mathcal{A} is closed under countable unions.

Proposition

Let \mathcal{A} be a nonempty algebra of sets of reals. Then

- \emptyset and $\mathbb{R} \in \mathcal{A}$.
- \mathcal{A} is closed under intersection.

Let \mathcal{A} be a nonempty σ -algebra of sets of reals. Then

- \mathcal{A} is closed under countable intersections.

MAT 5620: 3

Sigma Samples

Examples

1 $\mathcal{A} = \{\emptyset, \mathbb{R}\}$

2 $\mathcal{F} = \{F \subset \mathbb{R} : F \text{ is finite or } F^c \text{ is finite}\}$

1 \mathcal{F} is an algebra, the *co-finite algebra*

2 \mathcal{F} is not a σ -algebra

For each $r \in \mathbb{Q}$, the set $\{r\} \in \mathcal{F}$. But $\bigcup_{r \in \mathbb{Q}} \{r\} = \mathbb{Q} \notin \mathcal{F}$

3 Let $\mathcal{A} = \{\emptyset, [-1, 1], (-\infty, -1) \cup (1, \infty), \mathbb{R}\}$. Is \mathcal{A} an algebra?

4 Any intersection of σ -algebras is a σ -algebra

5 Let $\mathcal{B}(\mathbb{R})$ be the smallest σ -algebra containing all the open sets, the *Borel σ -algebra*.

MAT 5620: 4

Outer Measure

Definition (Lebesgue Outer Measure)

Let $E \subset \mathbb{R}$. Define the *Lebesgue Outer Measure* μ^* of E to be

$$\mu^*(E) = \inf_{E \subset \bigcup I_n} \sum_n \ell(I_n),$$

the infimum of the sums of the lengths of open interval covers of E .

Proposition (Monotonicity)

If $A \subseteq B$, then $\mu^*(A) \leq \mu^*(B)$.

Proposition

If I is an interval, then $\mu^*(I) = \ell(I)$.

MAT 5620: 5

Outer Measure of an Interval

Proof.

I. I is closed and bounded (compact). Then $I = [a, b]$.

- 1 For any $\varepsilon > 0$, $[a, b] \subset (a - \varepsilon, b + \varepsilon)$. So $\mu^*(I) \leq b - a + 2\varepsilon$. Since ε is arbitrary, $\mu^*(I) \leq b - a$.
- 2 Let $\{I_n\}$ cover $[a, b]$ with open intervals. There is a finite subcover for $[a, b]$. Order the subcover so that consecutive intervals overlap. Then

$$\sum_N \ell(I_k) = (b_1 - a_1) + (b_2 - a_2) + \cdots + (b_N - a_N)$$

Rearrange

$$\begin{aligned} \sum_N \ell(I_k) &= b_N - (a_N - b_{N-1}) - (a_{N-1} - b_{N-2}) - \cdots - (a_2 - b_1) - a_1 \\ &\geq b_N - a_1 > b - a \end{aligned}$$

Whence $\mu^*(I) = b - a$.

MAT 5620: 6

Outer Measure of an Interval, II

Proof (cont).

II. Let I be any bounded interval and $\varepsilon > 0$.

- ① There is a closed interval $J \subset I$ so that $\ell(I) - \varepsilon < \ell(J)$. Then

$$\ell(I) - \varepsilon < \ell(J) = \mu^*(J) \leq \mu^*(I) \leq \mu^*(\bar{I}) = \ell(\bar{I}) = \ell(I)$$

III. Suppose I is infinite.

- ① Then for each n , there is a closed interval $J \subset I$ s.t. $\ell(J) = n$
 ② Thence $\mu^*(I) \geq n$ for all n .

Aha! $\mu^*(I) = \infty$

Proposition

$$\mu^*(\mathbb{Q}) = 0$$

Proof.

Order \mathbb{Q} as $\{r_1, r_2, \dots\}$. $\{I_n = (r_n - \varepsilon/2^n, r_n + \varepsilon/2^n)\}$ covers \mathbb{Q} □

MAT 5620: 7

Countable Subadditivity

Theorem (μ^* is Countably Subadditive)

Let $\{E_n\}$ be a countable set sequence in \mathbb{R} . Then $\mu^*\left(\bigcup_n E_n\right) \leq \sum_n \mu^*(E_n)$

Proof.

I. If $\mu^*(E_n) = \infty$ for any n , then done.

II. Let $\varepsilon > 0$

- ① For each n find a cover $\{I_{n,j}\}_{j \in \mathbb{N}}$ such that $\sum_{j \in \mathbb{N}} \ell(I_{n,j}) < \mu^*(E_n) + \frac{\varepsilon}{2^n}$
 ② Then $\{I_{n,j}\}_{n,j \in \mathbb{N}}$ covers $E = \bigcup_n E_n$.

- ③ Whereupon

$$\begin{aligned} \mu^*(E) &\leq \sum_{n,j \in \mathbb{N}} \ell(I_{n,j}) = \sum_{n \in \mathbb{N}} \left[\sum_{j \in \mathbb{N}} \ell(I_{n,j}) \right] \\ &< \sum_{n \in \mathbb{N}} \left[\mu^*(E_n) + \frac{\varepsilon}{2^n} \right] = \sum_{n \in \mathbb{N}} [\mu^*(E_n)] + \varepsilon \end{aligned}$$

□

MAT 5620: 8

Open Holding & Lebesgue's Measure

Corollary

Given $E \subseteq \mathbb{R}$ and $\varepsilon > 0$, there is an open set $O \supseteq E$ s.t.

$$\mu^*(E) \leq \mu^*(O) \leq \mu^*(E) + \varepsilon$$

Definition (Carathéodory's Condition)

A set E is *Lebesgue measurable* iff for every (test) set A ,

$$\mu^*(A) = \mu^*(A \cap E) + \mu^*(A \cap E^c)$$

Let \mathfrak{M} be the collection of all Lebesgue measurable sets.

Corollary

For any A and E ,

$$\mu^*(A) = \mu^*((A \cap E) \cup (A \cap E^c)) \leq \mu^*(A \cap E) + \mu^*(A \cap E^c)$$

MAT 5620: 9

Much Ado About Nothing

Theorem

If $\mu^*(E) = 0$, then $E \in \mathfrak{M}$; i.e., E is measurable.

Proof.

Given the previous corollary, we need only show that

$$\mu^*(A \cap E) + \mu^*(A \cap E^c) \leq \mu^*(A)$$

- ① Since $A \cap E \subset E$, then $\mu^*(A \cap E) \leq \mu^*(E) = 0$.
- ② Since $A \cap E^c \subset A$, then $\mu^*(A \cap E^c) \leq \mu^*(A)$.

Whence $\mu^*(A \cap E) + \mu^*(A \cap E^c) \leq 0 + \mu^*(A) = \mu^*(A)$. □

Corollary

$$\mu^*(\mathbb{Q}) = 0 \implies \mathbb{Q} \in \mathfrak{M}$$

MAT 5620: 10

Unions Work

Theorem

A finite union of measurable sets is measurable.

Proof.

Let E_1 and $E_2 \in \mathfrak{M}$. Let A be a test set.

- 1 Use $A \cap E_1^c$ as a test set for E_2 which is measurable. Thence

$$\mu^*(A \cap E_1^c) = \mu^*((A \cap E_1^c) \cap E_2) + \mu^*((A \cap E_1^c) \cap E_2^c)$$

- 2 Note $A \cap (E_1 \cup E_2) = (A \cap E_1) \cup (A \cap E_2 \cap E_1^c)$. Whereupon

$$\begin{aligned} \mu^*(A \cap (E_1 \cup E_2)) + \mu^*(A \cap (E_1 \cup E_2)^c) &= \mu^*(A \cap (E_1 \cup E_2)) + \mu^*(A \cap (E_1^c \cap E_2^c)) \\ &\leq [\mu^*(A \cap E_1) + \mu^*(A \cap E_2 \cap E_1^c)] + \mu^*(A \cap E_1^c \cap E_2^c) \\ &\leq \mu^*(A \cap E_1) + [\mu^*(A \cap E_1^c \cap E_2) + \mu^*(A \cap E_1^c \cap E_2^c)] \\ &= \mu^*(A \cap E_1) + \mu^*(A \cap E_1^c) \\ &= \mu^*(A) \end{aligned}$$

□

Countable Unions Work

Theorem

The countable union of measurable sets is measurable.

Proof.

Let $E_k \in \mathfrak{M}$ and $E = \bigcup_n E_n$. Choose a test set A .

We need to show $\mu^*(A \cap E) + \mu^*(A \cap E^c) \leq \mu^*(A)$.

- 1 Set $F_n = \bigcup^n E_k$ and $F = \bigcup^\infty E_k = E$. Define $G_1 = E_1$, $G_2 = E_2 - E_1$, \dots , $G_k = E_k - \bigcup^{k-1} E_j$, and $G = \bigcup G_k$. Then

$$(i) G_i \cap G_j = \emptyset, (i \neq j) \quad (ii) F_n = \bigcup_n G_k \quad (iii) F = G = E$$

- 2 Test F_n with A to obtain $\mu^*(A) = \mu^*(A \cap F_n) + \mu^*(A \cap F_n^c)$

- 3 Test G_n with $A \cap F_n$ to obtain

$$\begin{aligned} \mu^*(A \cap F_n) &= \mu^*((A \cap F_n) \cap G_n) + \mu^*((A \cap F_n) \cap G_n^c) \\ &= \mu^*(A \cap G_n) + \mu^*(A \cap F_{n-1}) \end{aligned}$$

□

Countable Unions Work, II

Proof.

- ④ Iterate $\mu^*(A \cap F_n) = \mu^*(A \cap G_n) + \mu^*(A \cap F_{n-1})$ from 3 to have

$$\mu^*(A \cap F_n) = \sum_{k=1}^n \mu^*(A \cap G_k)$$

- ⑤ Since $F_n \subseteq F$, then $F^c \subseteq F_n^c$ for all n , then

$$\mu^*(A \cap F_n^c) \geq \mu^*(A \cap F^c)$$

- ⑥ Whence

$$\mu^*(A) \geq \sum_{k=1}^n \mu^*(A \cap G_k) + \mu^*(A \cap F^c)$$

The summation is increasing & bounded, so convergent.

- ⑦ However $\sum_{k=1}^{\infty} \mu^*(A \cap G_k) \geq \mu^*\left(\bigcup_{k=1}^{\infty} (A \cap G_k)\right) = \mu^*(A \cap F)$

Aha! $\mu^*(A) \geq \mu^*(A \cap F) + \mu^*(A \cap F^c)$ □

MAT 5620: 13

Everything Works

Corollary

The collection of Lebesgue measurable sets \mathfrak{M} is a σ -algebra.

Corollary

The Borel sets are measurable. (There are measurable, non-Borel sets.)

$$\mathcal{B}(\mathbb{R}) \subsetneq \mathfrak{M} \subsetneq \mathcal{P}(\mathbb{R})$$

Definition (Lebesgue Measure)

Lebesgue measure μ is μ^* restricted to \mathfrak{M} . So $\mu: \mathfrak{M} \rightarrow [0, \infty]$.

Definition (Almost Everywhere)

A property P holds *almost everywhere* (a.e.) iff $\mu(\{x : \neg P(x)\}) = 0$.

MAT 5620: 14

The Return of Additivity

Theorem

Let $\{E_n\}$ be a countable (finite or infinite) sequence of pairwise disjoint sets in \mathfrak{M} . Then

$$\mu\left(\bigcup_{k=1}^{\infty} E_k\right) = \sum_{k=1}^{\infty} \mu(E_k)$$

Proof.

I. n is finite.

- 1 For $n = 1$, \checkmark
- 2 $(\bigcup_{k=1}^n E_k) \cap E_n = E_n$ and $(\bigcup_{k=1}^n E_k) \cap E_n^c = \bigcup_{k=1}^{n-1} E_k$
- 3 $\mu(\bigcup_{k=1}^n E_k) = \mu([\bigcup_{k=1}^n E_k] \cap E_n) + \mu([\bigcup_{k=1}^n E_k] \cap E_n^c)$
 $= \mu(E_n) + \mu(\bigcup_{k=1}^{n-1} E_k) = \mu(E_n) + \sum_{k=1}^{n-1} \mu(E_k) = \sum_{k=1}^n \mu(E_k)$

II. n is infinite.

- 1 $\bigcup_{k=1}^n E_k \subset \bigcup_{k=1}^{\infty} E_k \implies \mu(\bigcup_{k=1}^n E_k) = \sum_{k=1}^n \mu(E_k) \leq \mu(\bigcup_{k=1}^{\infty} E_k)$
- 2 A bnded & incr sum converges. Thus $\sum_{k=1}^{\infty} \mu(E_k) \leq \mu(\bigcup_{k=1}^{\infty} E_k)$
- 3 Subadditivity finishes the proof. □

MAT 5620: 15

Adding an Example

Example

Set $E_n = \left(\frac{n-1}{n}, \frac{n}{n+1}\right)$ for $n = 1.. \infty$.

- 1 The E_n are pairwise disjoint.
- 2 $\mu(E_n) = \ell(E_n) = \frac{n}{n+1} - \frac{n-1}{n} = \frac{1}{n(n+1)}$
- 3 $\mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \sum_{n=1}^{\infty} \mu(E_n) = \sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \sum_{n=1}^{\infty} \left[\frac{1}{n} - \frac{1}{n+1}\right]$

Whence $\mu\left(\bigcup_{n=1}^{\infty} E_n\right) = 1$.

NOTA BENE: $\bigcup_{n=1}^{\infty} E_n = (0, 1) - \left\{\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \dots\right\}$. Hence $\bigcup_{n=1}^{\infty} E_n = (0, 1)$ a.e.

MAT 5620: 16

Matryoshka

Theorem

If $\{E_n\}$ is a seq of nested, measurable sets with $\mu(E_1) < \infty$, then

$$\mu\left(\bigcap_{n=1}^{\infty} E_n\right) = \lim_{n \rightarrow \infty} \mu(E_n)$$

Proof.

- 1 Set $E = \bigcap_{k=1}^{\infty} E_k$. Set $F_k = E_k - E_{k+1}$. The F_k are pairwise disjoint.
- 2 Since $\bigcup_{k=1}^{\infty} F_k = E_1 - E$, then $\mu(E_1 - E) = \sum_{k=1}^{\infty} \mu(F_k) = \sum_{k=1}^{\infty} \mu(E_k - E_{k+1})$.
- 3 If $A \supset B$, then $\mu(A - B) = \mu(A) - \mu(B)$. Apply to the formula above.
- 4 $\mu(E_1) - \mu(E) = \sum_{k=1}^{\infty} \mu(E_k) - \mu(E_{k+1}) = \mu(E_1) - \lim_{k \rightarrow \infty} \mu(E_k)$

Since $\mu(E_1)$ is finite, we're done. □

MAT 5620: 17

The Cantor Set

Cantor Sets¹

I. Constructing C

- 1 Set $C_0 = [0, 1]$
- 2 Set $C_1 = C_0 - (\frac{1}{3}, \frac{2}{3})$
- 3 Set $C_2 = C_1 - (\frac{1}{3^2}, \frac{2}{3^2}) - (\frac{7}{3^2}, \frac{8}{3^2})$
- 4 Set $C_3 = C_2 - (\frac{1}{3^3}, \frac{2}{3^3}) - (\frac{7}{3^3}, \frac{8}{3^3}) - (\frac{19}{3^3}, \frac{20}{3^3}) - (\frac{25}{3^3}, \frac{26}{3^3})$
- 5 Let $C = \bigcap C_i$

II. Properties of C

- | | |
|--|---|
| 1 $\mu(C_0) = 1, \mu(C_1) = 2/3,$
$\mu(C_2) = 4/9, \mu(C_3) = 8/27, \dots$ So
$\mu(C_n) = \frac{2}{3}\mu(C_{n-1}) = \frac{2^n}{3^n}$ Whence
$\mu(C) = 0.$ | 4 C is nowhere dense |
| 2 C is uncountable | 5 C is compact |
| 3 C is perfect | 6 C is totally disconnected |
| | 7 $(\forall i) \partial C_i \subset C$ |
| | 8 $(\forall i) \frac{1}{4} \notin \partial C_i, \text{ but } \frac{1}{4} \in C$ |

¹Cantor gave the set in a footnote to show "perfect" $\not\subset$ "everywhere dense".

MAT 5620: 18

Not So Strange After All

Theorem

Let $E \subseteq \mathbb{R}$ and let $\varepsilon > 0$. TFAE:

- ① E is measurable
- ② There is an open set $O \supset E$ s.t. $\mu^*(O - E) < \varepsilon$
- ③ There is a closed set $F \subset E$ s.t. $\mu^*(E - F) < \varepsilon$

Proposition

Let S and T be measurable subsets of \mathbb{R} . Then

$$\mu(S \cup T) + \mu(S \cap T) = \mu(S) + \mu(T)$$

MAT 5620: 19

Functionally Measurable

Theorem (Measurability Conditions for Functions)

Let $f: D \rightarrow \mathbb{R}_\infty$ for some $D \in \mathfrak{M}$. TFAE

- ① For each $r \in \mathbb{R}$, the set $f^{-1}((r, \infty))$ is measurable.
- ② For each $r \in \mathbb{R}$, the set $f^{-1}([r, \infty))$ is measurable.
- ③ For each $r \in \mathbb{R}$, the set $f^{-1}((-\infty, r))$ is measurable.
- ④ For each $r \in \mathbb{R}$, the set $f^{-1}((-\infty, r])$ is measurable.

Proof.

$$1 \Rightarrow 2: \{x \mid f(x) \geq r\} = \bigcap_n \{x \mid f(x) > r - 1/n\}$$

$$2 \Rightarrow 3: \{x \mid f(x) < r\} = D - \{x \mid f(x) \geq r\}$$

$$3 \Rightarrow 4: \{x \mid f(x) \leq r\} = \bigcap_n \{x \mid f(x) < r + 1/n\}$$

$$4 \Rightarrow 1: \{x \mid f(x) > r\} = D - \{x \mid f(x) \leq r\} \quad \square$$

WmCB 2020. 20

The Measurably Functional

Corollary

If f satisfies any measurability condition, then $\{x \mid f(x) = r\}$ is measurable for each r .

Definition (Measurable Function)

If a function $f: D \rightarrow \mathbb{R}_\infty$ has measurable domain D and satisfies any of the measurability conditions, then f is *measurable*.

Definition

Step function: $\phi: [a, b] \rightarrow \mathbb{R}_\infty$ is a *step function* if there is a partition $a = x_0 < x_1 < \dots < x_n = b$ s.t. ϕ is constant on each interval $I_k = (x_{k-1}, x_k)$, then

$$\phi(x) = \sum_{k=1}^n a_k \chi_{I_k}(x)$$

Simple function: A function ψ with range $\{a_1, a_2, \dots, a_n\}$ where each set $\psi^{-1}(a_k)$ is measurable is a *simple function*.

MAT 5620: 21

Simply Stepping

Proposition

Step functions and simple functions are measurable

Theorem (Algebra of Measurable Functions)

Let f and g be measurable on a common domain D , and let $c \in \mathbb{R}$. Then

① $f + c$

③ $f \pm g$

⑤ $f \cdot g$

② $c \cdot f$

④ f^2

are all measurable.

Proof.

• ✓

□

MAT 5620: 22

Sequencing

Theorem

Let $\{f_n\}$ be a sequence of measurable functions on a common domain D . Then

- | | | |
|------------------------------|-------------------------------------|--|
| 1 $\sup \{f_1, \dots, f_n\}$ | 3 $\sup_{n \rightarrow \infty} f_n$ | 5 $\limsup_{n \rightarrow \infty} f_n$ |
| 2 $\inf \{f_1, \dots, f_n\}$ | 4 $\inf_{n \rightarrow \infty} f_n$ | 6 $\liminf_{n \rightarrow \infty} f_n$ |

are all measurable.

Proof.

1 Set $f = \{f_1, \dots, f_n\}$. Then $\{f(x) > r\} = \bigcup_{k=1}^n \{f_k(x) > r\}$.

3 Set $F = \sup_n f_n$. Then $\{F(x) > r\} = \bigcup_{k=1}^{\infty} \{f_k(x) > r\}$.

5 Set $\Phi = \limsup_n f_n$. Then $\limsup_{n \rightarrow \infty} f_n = \inf_n \left[\sup_{k \geq n} f_k \right]$

□

MAT 5620: 23

Zeroing

Theorem

If f is measurable and $f = g$ a.e., then g is measurable.

Definition (Convergence Almost Everywhere)

A sequence $\{f_n\}$ converges to f almost everywhere, written as $f_n \rightarrow f$ a.e., iff $\mu(\{x: f_n(x) \not\rightarrow f(x)\}) = 0$.

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$. Then f is measurable iff there is a seq. of simple functions $\{\psi_n\}$ converging to f a.e.

MAT 5620: 24

A Simple Proof

Proof.

(\Rightarrow) Wolog $f \geq 0$.

① Define $A_{n,k} = \{x \mid \frac{k-1}{2^n} \leq f(x) < \frac{k}{2^n}\}$ for $k = 1..(n \cdot 2^n)$ and

$$A_{0,n} = [a, b] - \bigcup_{k=1}^{n2^n} A_{n,k}$$

② Set $\psi_n(x) = n\chi_{A_{0,n}}(x) + \sum_{k=1}^{n2^n} \frac{k-1}{2^n} \cdot \chi_{A_{n,k}}(x)$

③ Then

① $\psi_1 \leq \psi_2 \leq \dots$

② If $0 \leq f(x) \leq n$, then $|f - \psi_n| < 2^{-n}$

③ $\lim_n \psi = f$ a.e.

(\Leftarrow) ✓ □

MAT 5620: 25

Integration

We began by looking at two examples of integration problems.

- The Riemann integral over $[0, 1]$ of a function with infinitely many discontinuities didn't exist even though the points of discontinuity formed a set of measure zero.
(The points of discontinuity formed a dense set in $[0, 1]$.)
- The limit of a sequence of Riemann integrable functions did not equal the integral of the limit function of the sequence.
(Each function had area $1/2$, but the limit of the sequence was the zero function.)

We will look at Riemann integration, then Riemann-Stieltjes integration, and last, develop the Lebesgue integral.

There are many other types of integrals: Darboux, Denjoy, Gauge, Perron, etc. See the list given in the "See also" section of *Integrals* on Mathworld.

MAT 5620: 26

Riemann Integral

Definition

- A *partition* \mathcal{P} of $[a, b]$ is a finite set of points such that $\mathcal{P} = \{a = x_0 < x_1 < \cdots < x_{n-1} < x_n = b\}$.
- Set $M_i = \sup f(x)$ on $[x_{i-1}, x_i]$. The *upper sum* of f on $[a, b]$ w.r.t. \mathcal{P} is

$$U(\mathcal{P}, f) = \sum_{i=1}^n M_i \cdot \Delta x_i$$

- The *upper Riemann integral* of f over $[a, b]$ is

$$\int_a^b f(x) dx = \inf_{\mathcal{P}} U(\mathcal{P}, f)$$

Exercise

- 1 Define the lower sum $L(\mathcal{P}, f)$ and the lower integral $\int_a^b f$.

MAT 5620: 27

Definitely a Riemann Integral

Definition

If $\int_a^b f(x) dx = \int_a^b f(x) dx$, then f is Riemann integrable and is written as $\int_a^b f(x) dx$ and $f \in \mathfrak{R}$ on $[a, b]$.

Proposition

A function f is Riemann integrable on $[a, b]$ if and only if for every $\epsilon > 0$ there is a partition \mathcal{P} of $[a, b]$ such that

$$U(\mathcal{P}, f) - L(\mathcal{P}, f) < \epsilon.$$

Theorem

If f is continuous on $[a, b]$, then $f \in \mathfrak{R}$ on $[a, b]$.

Theorem

If f is bounded on $[a, b]$ with only finitely many points of discontinuity, then $f \in \mathfrak{R}$ on $[a, b]$.

MAT 5620: 28

Properties of Riemann Integrals

Proposition

Let f and $g \in \mathfrak{R}$ on $[a, b]$ and $c \in \mathbb{R}$. Then

- $\int_a^b cf \, dx = c \int_a^b f \, dx$
- $\int_a^b (f + g) \, dx = \int_a^b f \, dx + \int_a^b g \, dx$
- $f \cdot g \in \mathfrak{R}$
- if $f \leq g$, then $\int_a^b f \, dx \leq \int_a^b g \, dx$
- $\left| \int_a^b f \, dx \right| \leq \int_a^b |f| \, dx$
- Define $F(x) = \int_a^x f(t) \, dt$. Then F is continuous and, if f is continuous at x_0 , then $F'(x_0) = f(x_0)$
- If $F' = f$ on $[a, b]$, then $\int_a^b f(x) \, dx = F(b) - F(a)$

MAT 5620: 29

Riemann Integrated Exercises

Exercises

- 1 If $\int_a^b |f(x)| \, dx = 0$, then $f = 0$.
- 2 Show why $\int_0^1 \chi_{\mathbb{Q}}(x) \, dx$ does not exist.
- 3 Define

$$S_n(x) = \sum_{k=1}^{n+1} \left(\frac{k-1}{k} \cdot \chi_{\left[\frac{k-1}{k}, \frac{k}{k+1}\right)}(x) \right) + \frac{n}{n+1} \chi_{\left[\frac{n+1}{n+2}, 1\right]}(x).$$

- 1 How many discontinuities does S_n have?
- 2 Prove that $S'_n(x) = 0$ a.e.
- 3 Calculate $\int_0^1 S_n(x) \, dx$.
- 4 What is S_∞ ?
- 5 Does $\int_0^1 S_\infty(x) \, dx$ exist?

(See an animated graph of S_N .)

MAT 5620: 30

Riemann-Stieltjes Integral

Definition

- Let $\alpha(x)$ be a monotonically increasing function on $[a, b]$. Set $\Delta\alpha_i = \alpha(x_i) - \alpha(x_{i-1})$.
- Set $M_i = \sup f(x)$ on $[x_{i-1}, x_i]$. The *upper sum* of f on $[a, b]$ w.r.t. α and \mathcal{P} is

$$U(\mathcal{P}, f, \alpha) = \sum_{i=1}^n M_i \cdot \Delta\alpha_i$$

- The *upper Riemann-Stieltjes integral* of f over $[a, b]$ w.r.t. α is

$$\int_a^b f(x) d\alpha(x) = \inf_{\mathcal{P}} U(\mathcal{P}, f, \alpha)$$

Exercise

- 1 Define the lower sum $L(\mathcal{P}, f, \alpha)$ and lower integral $\int_a^b f d\alpha$.

MAT 5620: 31

Definitely a Riemann-Stieltjes Integral

Definition

If $\int_a^b f d\alpha = \int_a^b f d\alpha$, then f is Riemann-Stieltjes integrable and is written as $\int_a^b f(x) d\alpha(x)$ and $f \in \mathfrak{R}(\alpha)$ on $[a, b]$.

Proposition

A function f is Riemann-Stieltjes integrable w.r.t. α on $[a, b]$ iff for every $\epsilon > 0$ there is a partition \mathcal{P} of $[a, b]$ such that

$$U(\mathcal{P}, f, \alpha) - L(\mathcal{P}, f, \alpha) < \epsilon.$$

Theorem

If f is continuous on $[a, b]$, then $f \in \mathfrak{R}(\alpha)$ on $[a, b]$.

Theorem

If f is bounded on $[a, b]$ with only finitely many points of discontinuity and α is continuous at each of f 's discontinuities, then $f \in \mathfrak{R}(\alpha)$ on $[a, b]$.

MAT 5620: 32

Properties of Riemann-Stieltjes Integrals

Proposition

Let f and $g \in \mathfrak{R}(\alpha)$ and in β on $[a, b]$ and $c \in \mathbb{R}$. Then

- $\int_a^b cf \, d\alpha = c \int_a^b f \, d\alpha$ and $\int_a^b f \, d(c\alpha) = c \int_a^b f \, d\alpha$
- $\int_a^b (f + g) \, d\alpha = \int_a^b f \, d\alpha + \int_a^b g \, d\alpha$ and
 $\int_a^b f \, d(\alpha + \beta) = \int_a^b f \, d\alpha + \int_a^b f \, d\beta$
- $f \cdot g \in \mathfrak{R}(\alpha)$
- if $f \leq g$, then $\int_a^b f \, d\alpha \leq \int_a^b g \, d\alpha$
- $\left| \int_a^b f \, d\alpha \right| \leq \int_a^b |f| \, d\alpha$
- Suppose that $\alpha' \in \mathfrak{R}$ and f is bounded. Then $f \in \mathfrak{R}(\alpha)$ iff $f\alpha' \in \mathfrak{R}$ and

$$\int_a^b f \, d\alpha = \int_a^b f \cdot \alpha' \, dx$$

MAT 5620: 33

Riemann-Stieltjes Integrals and Series

Proposition

If f is continuous at $c \in (a, b)$ and $\alpha(x) = r$ for $a \leq x < c$ and $\alpha(x) = s$ for $c < x \leq b$, then

$$\begin{aligned} \int_a^b f \, d\alpha &= f(c) (\alpha(c+) - \alpha(c-)) \\ &= f(c) (s - r) \end{aligned}$$

Proposition

Let $\alpha = [x]$, the greatest integer function. If f is continuous on $[0, b]$, then

$$\int_0^b f(x) \, d[x] = \sum_{k=1}^{\lfloor b \rfloor} f(k)$$

MAT 5620: 34

Riemann-Stieltjes Integrated Exercises

Exercises

1 $\int_0^1 x dx^2$

2 $\int_0^{\pi/2} \cos(x) d \sin(x)$

3 $\int_0^{5/2} x d(x - \lfloor x \rfloor)$

4 $\int_{-1}^1 e^x d|x|$

5 $\int_{-3/2}^{3/2} e^x d\lfloor x \rfloor$

6 $\int_{-1}^1 e^x d\lfloor x \rfloor$

7 Set H to be the Heaviside function; i.e.,

$$H(x) = \begin{cases} 0 & x \leq 0 \\ 1 & \text{otherwise} \end{cases}.$$

Show that, if f is continuous at 0, then

$$\int_{-\infty}^{+\infty} f(x) dH(x) = f(0).$$

MAT 5620: 35

Lebesgue Integral

We start with **simple functions**.

Definition

A function has *finite support* if it vanishes outside a finite interval.

Definition

Let ϕ be a measurable simple function with finite support. If

$\phi(x) = \sum_{i=1}^n a_i \chi_{A_i}(x)$ is a representation of ϕ , then

$$\int \phi(x) dx = \sum_{i=1}^n a_i \cdot \mu(A_i)$$

Definition

If E is a measurable set, then $\int_E \phi = \int \phi \cdot \chi_E$.

Integral Linearity

Proposition

If ϕ and ψ are measurable simple functions with finite support and $a, b \in \mathbb{R}$, then $\int (a\phi + b\psi) = a \int \phi + b \int \psi$. Further, if $\phi \leq \psi$ a.e., then $\int \phi \leq \int \psi$.

Proof (sketch).

I. Let $\phi = \sum_{i=1}^N \alpha_i \chi_{A_i}$ and $\psi = \sum_{j=1}^M \beta_j \chi_{B_j}$. Then show $a\phi + b\psi$ can be written as $a\phi + b\psi = \sum_{k=1}^K (a\alpha_{k_i} + b\beta_{k_j}) \chi_{E_k}$ for the properly chosen E_k . Set A_0 and B_0 to be zero sets of ϕ and ψ . (Take $\{E_k : k = 0..K\} = \{A_j \cap B_k : j = 0..N, k = 0..M\}$.)

II. Use the definition to show $\int \psi - \int \phi = \int (\psi - \phi) \geq \int 0 = 0$. □

Steps to the Lebesgue Integral

Proposition

Let f be bounded on $E \in \mathfrak{M}$ with $\mu(E) < \infty$. Then f is measurable iff

$$\inf_{f \leq \psi} \int_E \psi = \sup_{f \geq \phi} \int_E \phi$$

for all simple functions ϕ and ψ .

Proof.

I. Suppose f is bounded by M . Define

$$E_k = \left\{ x : \frac{k-1}{n} M < f(x) \leq \frac{k}{n} M \right\}, \quad -n \leq k \leq n$$

The E_k are measurable, disjoint, and have union E . Set

$$\psi_n(x) = \frac{M}{n} \sum_{-n}^n k \chi_{E_k}(x), \quad \phi_n(x) = \frac{M}{n} \sum_{-n}^n (k-1) \chi_{E_k}(x)$$

□

SLI (cont)

(proof cont).

Then $\phi_n(x) \leq f(x) \leq \psi(x)$, and so

- $\inf \int_E \psi \leq \int_E \psi_n = \frac{M}{n} \sum_{k=-n}^n k \mu(E_k)$
- $\sup \int_E \phi \geq \int_E \phi_n = \frac{M}{n} \sum_{k=-n}^n (k-1) \mu(E_k)$


Thus $0 \leq \inf \int_E \psi - \sup \int_E \phi \leq \frac{M}{n} \mu(E)$. Since n is arbitrary, equality holds.

II. Suppose that $\inf \int_E \psi = \sup \int_E \phi$. Choose ϕ_n and ψ_n so that $\phi_n \leq f \leq \psi_n$ and $\int_E (\psi_n - \phi_n) < \frac{1}{n}$. The functions $\psi^* = \inf \psi_n$ and $\phi^* = \sup \phi_n$ are measurable and $\phi^* \leq f \leq \psi^*$. The set $\Delta = \{x : \phi^*(x) < \psi^*(x)\}$ has measure 0. Thus $\phi^* = \psi^*$ almost everywhere, so $\phi^* = f$ a.e. Hence f is measurable. \square

MAT 5620: 39

Example Steps

Example



Defining the Lebesgue Integral

Definition

If f is a bounded measurable function on a measurable set E with $m(E) < \infty$, then

$$\int_E f = \inf_{\psi \geq f} \int_E \psi$$

for all simple functions $\psi \geq f$.

Proposition

Let f be a bounded function defined on $E = [a, b]$. If f is Riemann integrable on $[a, b]$, then f is measurable on $[a, b]$ and

$$\int_E f = \int_a^b f(x) dx;$$

the Riemann integral of f equals the Lebesgue integral of f .

MAT 5620: 41

Properties of the Lebesgue Integral

Proposition

If f and g are measurable on E , a set of finite measure, then

- $\int_E (\alpha f + \beta g) = \alpha \int_E f + \beta \int_E g$
- if $f = g$ a.e., then $\int_E f = \int_E g$
- if $f \leq g$ a.e., then $\int_E f \leq \int_E g$
- $\left| \int_E f \right| \leq \int_E |f|$
- if $a \leq f \leq b$, then $a \cdot \mu(E) \leq \int_E f \leq b \cdot \mu(E)$
- if $A \cap B = \emptyset$, then $\int_{A \cup B} f = \int_A f + \int_B f$

MAT 5620: 42

Lebesgue Integral Examples

Examples

1 Let $T(x) = \begin{cases} \frac{1}{q} & x = \frac{p}{q} \in \mathbb{Q} \\ 0 & \text{otherwise} \end{cases}$. Then $\int_{[0,1]} T = \int_0^1 T(x) dx$.

2 Let $\chi_{\mathbb{Q}}(x) = \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & \text{otherwise} \end{cases}$. Then $\int_{[0,1]} \chi_{\mathbb{Q}} \neq \int_0^1 \chi_{\mathbb{Q}}(x) dx$.

3 Define

$$f_n(x) = \sum_{k=1}^{n+1} \left(\frac{k-1}{k} \cdot \chi_{\left[\frac{k-1}{k}, \frac{k}{k+1}\right)}(x) \right) + \frac{n}{n+1} \chi_{\left[\frac{n+1}{n+2}, 1\right]}(x).$$

Then

1 f_n is a step function, hence integrable

2 $f'_n(x) = 0$ a.e.

3 $\frac{1}{4} \leq \int_{[0,1]} f_n = \int_0^1 f_n(x) dx < \frac{3}{8}$

MAT 5620: 43

Extending the Integral Definition

Definition

Let f be a nonnegative measurable function defined on a measurable set E . Define

$$\int_E f = \sup_{h \leq f} \int_E h$$

where h is a bounded measurable function with finite support.

Proposition

If f and g are nonnegative measurable functions, then

- $\int_E c f = c \int_E f$ for $c > 0$
- $\int_E f + g = \int_E f + \int_E g$
- If $f \leq g$ a.e., then $\int_E f \leq \int_E g$

MAT 5620: 44

General Lebesgue's Integral

Definition

Set $f^+(x) = \max\{f(x), 0\}$ and $f^-(x) = \max\{-f(x), 0\}$. Then $f = f^+ - f^-$ and $|f| = f^+ + f^-$. A measurable function f is integrable over E iff both f^+ and f^- are integrable over E , and then $\int_E f = \int_E f^+ - \int_E f^-$.

Proposition

Let f and g be integrable over E and let $c \in \mathbb{R}$. Then

- 1 $\int_E cf = c \int_E f$
- 2 $\int_E f + g = \int_E f + \int_E g$
- 3 if $f \leq g$ a.e., then $\int_E f \leq \int_E g$
- 4 if A, B are disjoint m'ble subsets of E , $\int_{A \cup B} f = \int_A f + \int_B f$

MAT 5620: 45

Convergence Theorems

Theorem (Bounded Convergence Theorem)

Let $\{f_n : E \rightarrow \mathbb{R}\}$ be a sequence of measurable functions converging to f with $m(E) < \infty$. If there is a uniform bound M for all f_n , then

$$\int_E \lim_n f_n = \lim_n \int_E f_n$$

Proof (sketch).

Let $\epsilon > 0$.

- 1 f_n converges "almost uniformly;" i.e., $\exists A, N$ s.t. $m(A) < \frac{\epsilon}{4M}$ and, for $n > N$,
 $x \in E - A \implies |f_n(x) - f(x)| \leq \frac{\epsilon}{2m(E)}$.
- 2 $\left| \int_E f_n - \int_E f \right| = \left| \int_E f_n - f \right| \leq \int_E |f_n - f| = \left(\int_{E-A} + \int_A \right) |f_n - f|$
- 3 $\int_{E-A} |f_n - f| + \int_A |f_n| + |f| \leq \frac{\epsilon}{2m(E)} \cdot m(E) + 2M \cdot \frac{\epsilon}{4M} = \epsilon$

□

MAT 5620: 46

Lebesgue's Dominated Convergence Theorem

Theorem (Dominated Convergence Theorem)

Let $\{f_n : E \rightarrow \mathbb{R}\}$ be a sequence of measurable functions converging a.e. on E with $m(E) < \infty$. If there is an integrable function g on E such that $|f_n| \leq g$ then

$$\int_E \lim_n f_n = \lim_n \int_E f_n$$

Lemma

Under the conditions of the DCT, set $g_n = \sup_{k \geq n} \{f_n, f_{n+1}, \dots\}$ and

$h_n = \inf_{k \geq n} \{f_n, f_{n+1}, \dots\}$. Then g_n and h_n are integrable and $\lim g_n = f = \lim h_n$ a.e.

Proof of DCT (sketch).

- Both g_n and h_n are monotone and converging. Apply MCT.
- $h_n \leq f_n \leq g_n \implies \int_E h_n \leq \int_E f_n \leq \int_E g_n$. □

MAT 5620: 47

Increasing the Convergence

Theorem (Fatou's Lemma)

If $\{f_n\}$ is a sequence of measurable functions converging to f a.e. on E , then

$$\int_E \lim_n f_n \leq \lim_n \inf \int_E f_n$$

Theorem (Monotone Convergence Theorem)

If $\{f_n\}$ is an increasing sequence of nonnegative measurable functions converging to f , then

$$\int \lim_n f_n = \lim_n \int f_n$$

Corollary (Beppo Levi Theorem (cf.))

If $\{f_n\}$ is a sequence of nonnegative measurable functions, then

$$\int \sum_{n=1}^{\infty} f_n = \sum_{n=1}^{\infty} \int f_n$$

MAT 5620: 48

Sidebar: Littlewood's Three Principles

John Edensor Littlewood said,

The extent of knowledge required is nothing so great as sometimes supposed. There are three principles, roughly expressible in the following terms:

- *every measurable set is nearly a finite union of intervals;*
- *every measurable function is nearly continuous;*
- *every convergent sequence of measurable functions is nearly uniformly convergent.*

Most of the results of analysis are fairly intuitive applications of these ideas.

From *Lectures on the Theory of Functions*, Oxford, 1944, p. 26.

MAT 5620: 49

Extensions of Convergence

The sequence f_n converges to $f \dots$

Definition (Convergence Almost Everywhere)

almost everywhere if $m(\{x : f_n(x) \not\rightarrow f(x)\}) = 0$.

Definition (Convergence Almost Uniformly)

almost uniformly on E if, for any $\epsilon > 0$, there is a set $A \subset E$ with $m(A) < \epsilon$ so that f_n converges uniformly on $E - A$.

Definition (Convergence in Measure)

in measure if, for any $\epsilon > 0$, $\lim_{n \rightarrow \infty} m(\{x : |f_n(x) - f(x)| \geq \epsilon\}) = 0$.

Definition (Convergence in Mean (of order $p > 1$))

in mean if $\lim_{n \rightarrow \infty} \|f_n - f\|_p = \lim_{n \rightarrow \infty} \left[\int_E |f - f_n|^p \right]^{1/p} = 0$

MAT 5620: 50

Integrated Exercises

Exercises

- ① Prove: If f is integrable on E , then $|f|$ is integrable on E .
- ② Prove: If f is integrable over E , then $\left| \int_E f \right| \leq \int_E |f|$.
- ③ True or False: If $|f|$ is integrable over E , then f is integrable over E .
- ④ Let f be integrable over E . For any $\epsilon > 0$, there is a simple (resp. step) function ϕ (resp. ψ) such that $\int_E |f - \phi| < \epsilon$.
- ⑤ For $n = k + 2^\nu$, $0 \leq k < 2^\nu$, define $f_n = \chi_{[k2^{-\nu}, (k+1)2^{-\nu}]}$.
 - ① Show that f_n does not converge for any $x \in [0, 1]$.
 - ② Show that f_n does not converge a.e. on $[0, 1]$.
 - ③ Show that f_n does not converge almost uniformly on $[0, 1]$.
 - ④ Show that $f_n \rightarrow 0$ in measure.
 - ⑤ Show that $f_n \rightarrow 0$ in mean (of order 2).

MAT 5620: 51

References

Texts on analysis, integration, and measure:

- *Mathematical Analysis*, T. Apostle
- *Principles of Mathematical Analysis*, W. Rudin
- *Real Analysis*, H. Royden
- *Lebesgue Integration*, S. Chae
- *Geometric Measure Theory*, F. Morgan

Comparison of different types of integrals:

- *A Garden of Integrals*, F. Burk
- *Integral, Measure, and Derivative: A Unified Approach*, G. Shilov and B. Gurevich

MAT 5620: 52