


# MAT 5620. Analysis II. Notes on Measure Theory.

Wm C Bauldry

BauldryWC@appstate.edu

Autumn, 2006



Henri Léon Lebesgue  
(1875–1941)

# Table of Contents

- 0. “Riemann, We Have a Problem.”
- 1. Toward a Unit of Measure
  - 1.1 Set Algebras
    - ▶ Sidebar: Borel Sets
  - 1.2 Countably Additive Measures
  - 1.3 Outer Measure
  - 1.4 Measured Exercises
- 2. Lebesgue Measure
  - 2.1 Measurable Sets
  - 2.2 Measure Zero
    - ▶ Sidebar:  $\mathbb{Q}$  Is Small
  - 2.3 Measurable Functions
  - 2.4 Functionally Measured Exercises
- 3. Integration
  - 3.1 Riemann Integral
  - 3.2 Riemann-Stieltjes Integral
  - 3.3 Lebesgue Integral
  - 3.4 Convergence Theorems
    - ▶ Sidebar: Littlewood’s Three Principles
  - 3.5 Integrated Exercises
- 4. References

# “Riemann, We Have a Problem.”

There are problems with Riemann integration.

1. Define Dirichlet's function (1829)  $D(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{otherwise} \end{cases}$ .

Then  $\int_{[0,1]} D(x) dx$  does not exist.

2. Set  $f_n(x) = \begin{cases} 2n^2x & 0 \leq x < \frac{1}{2n} \\ 2n(1 - nx) & \frac{1}{2n} \leq x < \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$ . Then

$$\int_{[0,1]} \lim_{n \rightarrow \infty} f_n(x) dx \neq \lim_{n \rightarrow \infty} \int_{[0,1]} f_n(x) dx.$$

# “Riemann, We Have a Problem.”

There are problems with Riemann integration.

1. Define Dirichlet's function (1829)  $D(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{otherwise} \end{cases}$ .

Then  $\int_{[0,1]} D(x) dx$  does not exist.


2. Set  $f_n(x) = \begin{cases} 2n^2x & 0 \leq x < \frac{1}{2n} \\ 2n(1 - nx) & \frac{1}{2n} \leq x < \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$ . Then

$$\int_{[0,1]} \lim_{n \rightarrow \infty} f_n(x) dx \neq \lim_{n \rightarrow \infty} \int_{[0,1]} f_n(x) dx.$$

*Enter Henri Lebesgue in 1902.*

# A Bad Sequence of Functions

## Example



- Find  $\int f_n$ ,  $\lim_n \int f_n$ ,  $\lim_n f_n$ , and  $\int \lim_n f_n$ .

# Toward a Unit of Measure

## Definition

The *length of an interval* in  $\mathbb{R}^1$  is the difference of the endpoints and is given by  $\ell([a, b]) = b - a$ .

**Goal:** To have a set-function  $m : \mathfrak{M} \rightarrow \mathbb{R}$  that “measures” the “size” of a set where  $m$  ideally satisfies:

1.  $\mathfrak{M} = \mathcal{P}(\mathbb{R})$ ; *id est*, every set can be measured.
2. For every interval  $I$ , open or closed or not,  $m(I) = \ell(I)$ .
3. If the sequence  $\{E_n\}$  is disjoint, then  $m(\bigcup E_n) = \sum m(E_n)$ .
4.  $m$  is *translation invariant*; i.e.,  $m(E + x) = m(E)$  for every  $E$  and any  $x \in \mathbb{R}$ .

Unfortunately, this is impossible.<sup>1</sup> We give up the first and allow sets not to be in the class of measurable sets,  $\mathfrak{M} \subset \mathcal{P}(\mathbb{R})$ .

---

<sup>1</sup>Even the first 3 are impossible assuming the *continuum hypothesis*.

# $\sigma$ -Algebra of Sets

## Definition

Let  $\mathcal{A}$  be a collection of sets. Then  $\mathcal{A}$  is an *algebra of sets* or a *Boolean algebra* iff

- ▶ if  $A \in \mathcal{A}$ , then  $A^c \in \mathcal{A}$ ,
- ▶ if  $A, B \in \mathcal{A}$ , then  $A \cup B \in \mathcal{A}$ .

De Morgan's laws imply that if  $A, B \in \mathcal{A}$ , then  $A \cap B \in \mathcal{A}$ . Then we also have  $\emptyset \in \mathcal{A}$  and  $X \in \mathcal{A}$ .

## Definition

Let  $\mathcal{A}$  be an algebra of sets. Then  $\mathcal{A}$  is a  $\sigma$ -*algebra* of sets or *Borel field* iff for every countable sequence  $\{A_i\}$  of sets from  $\mathcal{A}$ , we have  $\bigcup A_i \in \mathcal{A}$ .

De Morgan's laws imply that countable intersections stay in  $\mathcal{A}$ .



# $\sigma$ -Algebra of Sets

## Definition

Let  $\mathcal{A}$  be a collection of sets. Then  $\mathcal{A}$  is an *algebra of sets* or a *Boolean algebra* iff

- ▶ if  $A \in \mathcal{A}$ , then  $A^c \in \mathcal{A}$ ,
- ▶ if  $A, B \in \mathcal{A}$ , then  $A \cup B \in \mathcal{A}$ .

De Morgan's laws imply that if  $A, B \in \mathcal{A}$ , then  $A \cap B \in \mathcal{A}$ . Then we also have  $\emptyset \in \mathcal{A}$  and  $X \in \mathcal{A}$ .

## Definition

Let  $\mathcal{A}$  be an algebra of sets. Then  $\mathcal{A}$  is a  $\sigma$ -*algebra* of sets or *Borel field* iff for every countable sequence  $\{A_i\}$  of sets from  $\mathcal{A}$ , we have  $\bigcup A_i \in \mathcal{A}$ .

De Morgan's laws imply that countable intersections stay in  $\mathcal{A}$ .

## Theorem

*There is a smallest  $\sigma$ -algebra containing any collection of sets.*

# Sidebar: Borel Sets

## Definition

The *Borel  $\sigma$ -algebra* on  $\mathbb{R}$  is the smallest  $\sigma$ -algebra containing  $\mathcal{G}$ , all of the open sets in  $\mathbb{R}$ , and is denoted by  $\mathcal{B}(\mathbb{R})$ .

## Proposition

The Borel  $\sigma$ -algebra  $\mathcal{B}(\mathbb{R})$  is (also) generated by each of:

- ▶  $\mathcal{F} = \{\text{all closed sets in } \mathbb{R}\}$
- ▶  $\{(-\infty, b] : b \in \mathbb{R}\}$
- ▶  $\{(a, b] : a, b \in \mathbb{R}\}$

## Proposition

Let  $\mathcal{S}_\delta = \{\bigcap S_i : S_i \in \mathcal{S}\}$  and  $\mathcal{S}_\sigma = \{\bigcup S_i : S_i \in \mathcal{S}\}$ . Then

$$\begin{array}{ccccccc} \mathcal{G} & \subsetneq & \mathcal{G}_\delta & \subsetneq & \mathcal{G}_{\delta\sigma} & \subsetneq & \mathcal{G}_{\delta\sigma\delta} \subsetneq \cdots \\ \circlearrowleft & & \circlearrowleft & & \circlearrowleft & & \circlearrowleft \quad \cdots \subsetneq \mathcal{B}(\mathbb{R}) \subsetneq \mathcal{P}(\mathbb{R}) \\ \mathcal{F} & \subsetneq & \mathcal{F}_\sigma & \subsetneq & \mathcal{F}_{\sigma\delta} & \subsetneq & \mathcal{F}_{\sigma\delta\sigma} \subsetneq \cdots \end{array}$$

# Countably Additive Measure

## Definition

A *countably additive measure* is a set function  $m$  such that

- ▶  $m$  is a non-negative extended real-valued function on a  $\sigma$ -algebra  $\mathfrak{M}$  of subsets of  $\mathbb{R}$ ; that is,  $m : \mathfrak{M} \rightarrow [0, \infty]$ .
- ▶  $m(\bigcup E_n) = \sum m(E_n)$  for any sequence of disjoint subsets.

# Countably Additive Measure

## Definition

A *countably additive measure* is a set function  $m$  such that

- ▶  $m$  is a non-negative extended real-valued function on a  $\sigma$ -algebra  $\mathfrak{M}$  of subsets of  $\mathbb{R}$ ; that is,  $m : \mathfrak{M} \rightarrow [0, \infty]$ .
- ▶  $m(\bigcup E_n) = \sum m(E_n)$  for any sequence of disjoint subsets.

## Exercises

Let  $m$  be a countably additive measure on the  $\sigma$ -algebra  $\mathfrak{M}$ .

1. If  $A$  and  $B$  are in  $\mathfrak{M}$  with  $A \subset B$ , then  $m(A) \leq m(B)$ .
2. If there is a set  $A \in \mathfrak{M}$  with  $m(A) < \infty$ , then  $m(\emptyset) = 0$ .
3. Show that  $m$  is countably subadditive or that for any sequence of sets,  $m(\bigcup E_n) \leq \sum m(E_n)$ . (Hint:  $B_n = A_n - \bigcup_{i < n} A_i$ .)
4. Let  $n$  be the counting measure, the number of elements in a set. Show that  $n$  satisfies Goals 1, 3, and 4.

# Outer Measure

## Definition

The *outer measure* of  $A$  is

$$m^*(A) = \inf_{A \subset \bigcup I_n} \sum_n \ell(I_n)$$

where  $I_n$  is open and  $\bigcup I_n$  covers  $A$  with a countable union.

## Proposition

*The outer measure of an interval is its length or  $m^*(I) = \ell(I)$ .*

# Outer Measure

## Definition

The *outer measure* of  $A$  is

$$m^*(A) = \inf_{A \subset \bigcup I_n} \sum_n \ell(I_n)$$

where  $I_n$  is open and  $\bigcup I_n$  covers  $A$  with a countable union.

## Proposition

*The outer measure of an interval is its length or  $m^*(I) = \ell(I)$ .*

## Proof.

- I.  $I = [a, b]$ . (a) Since  $[a, b] \subset (a - \epsilon, b + \epsilon)$ , then  $m^*(I) \leq b - a$ .  
(b) Heine-Borel thm: we need only consider finite covers. Work with the finite cover to show  $\sum \ell(I_n) \geq b - a$ .

# Outer Measure

## Definition

The *outer measure* of  $A$  is

$$m^*(A) = \inf_{A \subset \bigcup I_n} \sum_n \ell(I_n)$$

where  $I_n$  is open and  $\bigcup I_n$  covers  $A$  with a countable union.

## Proposition

*The outer measure of an interval is its length or  $m^*(I) = \ell(I)$ .*

## Proof.

- I.  $I = [a, b]$ . (a) Since  $[a, b] \subset (a - \epsilon, b + \epsilon)$ , then  $m^*(I) \leq b - a$ .  
(b) Heine-Borel thm: we need only consider finite covers. Work with the finite cover to show  $\sum \ell(I_n) \geq b - a$ .
- II. Any finite interval  $I$ . There is a closed interval  $J \subset I$  such that  $\ell(I) - \epsilon \leq \ell(J) = m^*(J) \leq m^*(I) \leq m^*(\bar{I}) = \ell(I)$ .

# Outer Measure

## Definition

The *outer measure* of  $A$  is

$$m^*(A) = \inf_{A \subset \bigcup I_n} \sum_n \ell(I_n)$$

where  $I_n$  is open and  $\bigcup I_n$  covers  $A$  with a countable union.

## Proposition

The outer measure of an interval is its length or  $m^*(I) = \ell(I)$ .

## Proof.

- I.  $I = [a, b]$ . (a) Since  $[a, b] \subset (a - \epsilon, b + \epsilon)$ , then  $m^*(I) \leq b - a$ .
- (b) Heine-Borel thm: we need only consider finite covers. Work with the finite cover to show  $\sum \ell(I_n) \geq b - a$ .
- II. Any finite interval  $I$ . There is a closed interval  $J \subset I$  such that  $\ell(I) - \epsilon \leq \ell(J) = m^*(J) \leq m^*(I) \leq m^*(\bar{I}) = \ell(I)$ .
- III. Any infinite interval. ✓





# Outer Measure is Countably Subadditive

## Theorem

Let  $\{A_n\}$  be a countable collection of subsets of  $\mathbb{R}$ . Then

$$m^* \left( \bigcup_n A_n \right) \leq \sum_n m^*(A_n)$$

# Outer Measure is Countably Subadditive

## Theorem

Let  $\{A_n\}$  be a countable collection of subsets of  $\mathbb{R}$ . Then

$$m^* \left( \bigcup_n A_n \right) \leq \sum_n m^*(A_n)$$

## Proof.

“Wolog” all  $A_n$ ’s have finite outer measure. For each  $A_n$  there is a countable collection of open intervals  $\{I_{n,i}\}$  covering  $A_n$  such that

$$\sum_i \ell(I_{n,i}) < m^*(A_n) + \frac{\epsilon}{2^n}$$

# Outer Measure is Countably Subadditive

## Theorem

Let  $\{A_n\}$  be a countable collection of subsets of  $\mathbb{R}$ . Then

$$m^* \left( \bigcup_n A_n \right) \leq \sum_n m^*(A_n)$$

## Proof.

“Wolog” all  $A_n$ ’s have finite outer measure. For each  $A_n$  there is a countable collection of open intervals  $\{I_{n,i}\}$  covering  $A_n$  such that

$$\sum_i \ell(I_{n,i}) < m^*(A_n) + \frac{\epsilon}{2^n}$$

The set  $\{I_{n,i} : n, i \in \mathbb{N}\}$  covers  $\bigcup A_n$ . Thence

$$m^* \left( \bigcup_n A_n \right) \leq \sum_{n,i} \ell(I_{n,i}) = \sum_n \sum_i \ell(I_{n,i}) < \sum_n \left( m^*(A_n) + \frac{\epsilon}{2^n} \right)$$



# Measured Exercises

## Exercises

1. *If  $A$  is a countable set, then  $m^*(A) = 0$ .*
2. *The closed interval  $[0, 1]$  is not countable.*
3. *Show that  $m^*(\mathbb{Q} \cap [0, 1]) = 0$  and  $m^*(\mathbb{Q}) = 0$ .*
4. *Let  $A = \mathbb{Q} \cap [0, 1]$  and let  $\{I_n : n = 1..N\}$  be a finite collection of open intervals covering  $A$ . Then  $\sum \ell(I_n) \geq 1$ .*
5. *Reconcile 1. through 4.*
6. *Given any set  $A$  and any  $\epsilon > 0$ , there is an open set  $G$  such that  $A \subset G$  and  $m^*(G) \leq m^*(A) + \epsilon$ .  
(Confer “Littlewood’s Three Principles.”)*
7. *Why is  $m^*$  translation invariant?*

# Lebesgue Measure

Lebesgue outer measure  $m^*$  satisfies [goals](#) 1, 2, and 4, but not goal 3, countable additivity;  $m^*$  is only countably subadditive. We can gain countable additivity by giving up goal 1 and reducing the collection  $\mathfrak{M}$  of sets; there will be sets that can't be measured. This approach is not without difficulties, though. The existence of nonmeasurable sets<sup>2</sup> leads to problems such as Vitali's theorem which yields a method of decomposing the interval  $[0, 1]$  into a set of measure 2. (Also see the Hausdorff paradox.)

We will use the definition of a set being measurable that was given by Carathéodory.

---

<sup>2</sup>See “Non-measurable\_set” for an intuitive explanation.

# Measurable Sets

## Definition

The set  $E$  is *measurable* iff for each set  $A$  we have

$$m^*(A) = m^*(A \cap E) + m^*(A \cap E^c).$$

## Proposition

If  $E$  is measurable, then  $E^c$  is measurable.

## Proposition

If  $m^*(E) = 0$ , then  $E$  is measurable.

## Proof.

Let  $A$  be any set. Then  $A \cap E \subset E$  implies  $m^*(A \cap E) \leq m^*(E)$ .

Hence  $m^*(A \cap E) = 0$ . Now  $A \cap E^c \subset A$ , so

$$m^*(A) \geq m^*(A \cap E^c) = m^*(A \cap E) + m^*(A \cap E^c). \quad \square$$

# Properties of “Measurable”

## Proposition

If  $E_1$  and  $E_2$  are measurable, then so is  $E_1 \cup E_2$ .

## Proof.

Let  $A$  be any set. Since  $E_2 \in \mathfrak{M}$ , then

$$m^*(A \cap E_1^c) = m^*((A \cap E_1^c) \cap E_2) + m^*((A \cap E_1^c) \cap E_2^c).$$

From  $A \cap (E_1 \cup E_2) = (A \cap E_1) \cup (A \cap E_2 \cap E_1^c)$ , we see that

$$m^*(A \cap (E_1 \cup E_2)) \leq m^*(A \cap E_1) + m^*(A \cap E_2 \cap E_1^c)$$

So

$$\begin{aligned} m^*(A \cap (E_1 \cup E_2)) + m^*(A \cap (E_1 \cup E_2)^c) & \\ & \leq m^*(A \cap E_1) + m^*(A \cap E_2 \cap E_1^c) + m^*(A \cap (E_1 \cup E_2)^c) \\ & = m^*(A \cap E_1) + m^*(A \cap E_1^c) = m^*(A) \end{aligned}$$



# Properties of “Measurable”

## Proposition

If  $E_1$  and  $E_2$  are measurable, then so is  $E_1 \cup E_2$ .

## Proof.

Let  $A$  be any set. Since  $E_2 \in \mathfrak{M}$ , then

$$m^*(A \cap E_1^c) = m^*((A \cap E_1^c) \cap E_2) + m^*((A \cap E_1^c) \cap E_2^c).$$

From  $A \cap (E_1 \cup E_2) = (A \cap E_1) \cup (A \cap E_2 \cap E_1^c)$ , we see that

$$m^*(A \cap (E_1 \cup E_2)) \leq m^*(A \cap E_1) + m^*(A \cap E_2 \cap E_1^c)$$

So

$$\begin{aligned} m^*(A \cap (E_1 \cup E_2)) + m^*(A \cap (E_1 \cup E_2)^c) & \\ & \leq m^*(A \cap E_1) + m^*(A \cap E_2 \cap E_1^c) + m^*(A \cap (E_1 \cup E_2)^c) \\ & = m^*(A \cap E_1) + m^*(A \cap E_1^c) = m^*(A) \end{aligned}$$

## Proposition



$\mathfrak{M}$  is an algebra of sets.



# A Bigger “Measurable” Cup

## Proposition

Let  $A$  be any set and  $E_1, E_2, \dots, E_N$  be a finite sequence of disjoint measurable sets. Then

$$m^* \left( A \cap \left[ \bigcup_{i=1}^N E_i \right] \right) = \sum_{i=1}^N m^*(A \cap E_i)$$

## Proof.

Induction on  $n$  with  $\left( A \cap \bigcup_{i=1}^n E_i \right) \cap E_n = A \cap E_n$  and

$$\left( A \cap \bigcup_{i=1}^n E_i \right) \cap E_n^c = A \cap \left( \bigcup_{i=1}^{n-1} E_i \right).$$

□

# A Countable “Measurable” Cup

## Proposition

Let  $E_1, E_2, \dots$  be a countable sequence of measurable sets. Then  $E = \bigcup_{i=1}^{\infty} E_i$  is measurable.

## Proof.

Wolog the  $E_i$  are pairwise disjoint. (Otherwise define  $B_i = E_i - \bigcup_{j=1}^{i-1} E_j$ .) Let  $A$  be any set and set  $F_n = \bigcup_{i=1}^n E_i$ . Then  $F_n \in \mathfrak{M}$  and  $F_n^c \supset E^c$ . Then  $m^*(A \cap F_n) = \sum_{i=1}^n m^*(A \cap E_i)$ . Hence, since  $n$  is arbitrary and  $m^*(A)$  is independent of  $n$ ,

$$m^*(A) \geq \sum_{i=1}^n m^*(A \cap E_i) + m^*(A \cap E^c) \geq m^*(A \cap E) + m^*(A \cap E^c)$$



# A Countable “Measurable” Cup

## Proposition

Let  $E_1, E_2, \dots$  be a countable sequence of measurable sets. Then  $E = \bigcup_{i=1}^{\infty} E_i$  is measurable.

## Proof.

Wolog the  $E_i$  are pairwise disjoint. (Otherwise define  $B_i = E_i - \bigcup_{j=1}^{i-1} E_j$ .) Let  $A$  be any set and set  $F_n = \bigcup_{i=1}^n E_i$ . Then  $F_n \in \mathfrak{M}$  and  $F_n^c \supset E^c$ . Then  $m^*(A \cap F_n) = \sum_{i=1}^n m^*(A \cap E_i)$ . Hence, since  $n$  is arbitrary and  $m^*(A)$  is independent of  $n$ ,

$$m^*(A) \geq \sum_{i=1}^n m^*(A \cap E_i) + m^*(A \cap E^c) \geq m^*(A \cap E) + m^*(A \cap E^c)$$

## Proposition

$\mathfrak{M}$  is a  $\sigma$ -algebra of sets.



# The Lebesgue Measure $m$ .

## Definition

Define Lebesgue measure to be the restriction  $m = m^*|_{\mathfrak{M}}$ .

# The Lebesgue Measure $m$ .

## Definition

Define Lebesgue measure to be the restriction  $m = m^*|_{\mathfrak{M}}$ .

## Theorem

*The Borel sets are Lebesgue measurable.*

# The Lebesgue Measure $m$ .

## Definition

Define Lebesgue measure to be the restriction  $m = m^*|_{\mathfrak{M}}$ .

## Theorem

*The Borel sets are Lebesgue measurable.*

## Theorem

Let  $E$  be a set and let  $\epsilon > 0$ . TFAE:

1.  $E$  is Lebesgue measurable
2. there is an open set  $G \supset E$  such that  $m^*(G - E) < \epsilon$
3. there is a closed set  $F \subset E$  such that  $m^*(E - F) < \epsilon$
4. there is a  $G \in \mathcal{G}_\delta$  such that  $E \subset G$  and  $m^*(G - E) = 0$
5. there is an  $F \in \mathcal{F}_\sigma$  such that  $F \subset E$  and  $m^*(E - F) = 0$

# Measure Zero

## Definition

A set  $S \subset \mathbb{R}$  has *measure zero* if and only if  $m(S) = 0$ ; i.e., for any  $\epsilon > 0$  there is an open cover  $\mathcal{C} = \{G_k \mid k \in \mathbb{N}\}$  of  $S$  such that  $\sum_{k \in \mathbb{N}} m(G_k) < \epsilon$ .

## Example

1. Any finite set (countable set) has measure zero.
2. Every interval  $[a, b]$  is not measure zero (when  $a < b$ ).

The length of  $[0, 1]$  is 1. The rationals contained in  $[0, 1]$  have measure zero. What is the measure of the irrationals in  $[0, 1]$ ?

## Definition (A.E.)

A property that holds for all  $x$  except on a set of measure zero is said to hold *almost everywhere*.

## Sidebar: $\mathbb{Q}$ Is Small

### Theorem

*The rationals are countable.*

### Proof.

*Let  $\mathbb{Q}$  be the set of rational numbers. The array below shows a method of enumerating all elements of  $\mathbb{Q}$ .*

$$\begin{array}{cccccc} 1/1_{(1)} & 2/1_{(2)} & 3/1_{(4)} & 4/1_{(7)} & \dots & \\ 1/2_{(3)} & 2/2_{(5)} & 3/2_{(8)} & 4/2_{(12)} & \dots & \\ 1/3_{(6)} & 2/3_{(9)} & 3/3_{(13)} & 4/3_{(18)} & \dots & \\ 1/4_{(10)} & 2/4_{(14)} & 3/4_{(19)} & 4/4_{(25)} & \dots & \\ \vdots & \vdots & \vdots & \vdots & \ddots & \end{array}$$

*Since each rational is counted, we have  $|\mathbb{Q}| \leq |\mathbb{N}|$  where we use  $|\cdot|$  to indicate cardinality (or size). But we know that  $\mathbb{N} \subseteq \mathbb{Q}$ , so that  $|\mathbb{N}| \leq |\mathbb{Q}|$ . Hence  $|\mathbb{Q}| = |\mathbb{N}|$ .*





# Covering $\mathbb{Q}$

## Theorem

*The set of rationals has measure zero.*

## Proof.

*Let  $\epsilon > 0$ . List the rationals in order  $\mathbb{Q} = \{r_1, r_2, r_3, \dots\}$  as given by the “countability matrix” defined earlier. For each rational  $r_k$ , define the open interval  $I_k = (r_k - \epsilon/2^{k+1}, r_k + \epsilon/2^{k+1})$ . Then*

- ▶ *the collection  $\mathcal{C} = \{I_k \mid k \in \mathbb{N}\}$  forms an open cover of  $\mathbb{Q}$ ,*
- ▶ *the length of each  $I_k$  is  $m(I_k) = \epsilon/2^k$ .*

*Then  $m(\mathbb{Q}) \leq m(\mathcal{C})$  which is*

$$m(\mathbb{Q}) \leq m(\mathcal{C}) = \sum_{k=1}^{\infty} m(I_k) = \sum_{k=1}^{\infty} \epsilon/2^k = \epsilon \sum_{k=1}^{\infty} \frac{1}{2^k} = \epsilon$$



# Measurable Functions

## Proposition (measurability condition)

Let  $f$  be an extended real-valued function on a measurable domain  $D$ . Then TFAE:

1. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) > \alpha\}$  is measurable.
2. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \geq \alpha\}$  is measurable.
3. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) < \alpha\}$  is measurable.
4. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \leq \alpha\}$  is measurable.

These imply

5. For each  $\beta \in \mathbb{R}_\infty$ , the set  $\{x : f(x) = \beta\}$  is measurable.

# Measurable Functions

## Proposition (measurability condition)

Let  $f$  be an extended real-valued function on a measurable domain  $D$ . Then TFAE:

1. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) > \alpha\}$  is measurable.
2. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \geq \alpha\}$  is measurable.
3. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) < \alpha\}$  is measurable.
4. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \leq \alpha\}$  is measurable.

These imply

5. For each  $\beta \in \mathbb{R}_\infty$ , the set  $\{x : f(x) = \beta\}$  is measurable.

## Proof.

$$(1.) \implies (2.) \quad \{x : f(x) \geq \alpha\} = \bigcap \{x : f(x) > \alpha - 1/n\}$$

# Measurable Functions

## Proposition (measurability condition)

Let  $f$  be an extended real-valued function on a measurable domain  $D$ . Then TFAE:

1. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) > \alpha\}$  is measurable.
2. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \geq \alpha\}$  is measurable.
3. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) < \alpha\}$  is measurable.
4. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \leq \alpha\}$  is measurable.

These imply

5. For each  $\beta \in \mathbb{R}_\infty$ , the set  $\{x : f(x) = \beta\}$  is measurable.

## Proof.

$$(1.) \implies (2.) \quad \{x : f(x) \geq \alpha\} = \bigcap \{x : f(x) > \alpha - 1/n\}$$

$$(2.) \implies (3.) \quad \{x : f(x) < \alpha\} = D - \{x : f(x) \geq \alpha\}$$

# Measurable Functions

## Proposition (measurability condition)

Let  $f$  be an extended real-valued function on a measurable domain  $D$ . Then TFAE:

1. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) > \alpha\}$  is measurable.
2. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \geq \alpha\}$  is measurable.
3. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) < \alpha\}$  is measurable.
4. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \leq \alpha\}$  is measurable.

These imply

5. For each  $\beta \in \mathbb{R}_\infty$ , the set  $\{x : f(x) = \beta\}$  is measurable.

## Proof.

$$(1.) \implies (2.) \quad \{x : f(x) \geq \alpha\} = \bigcap \{x : f(x) > \alpha - 1/n\}$$

$$(2.) \implies (3.) \quad \{x : f(x) < \alpha\} = D - \{x : f(x) \geq \alpha\}$$

$$(3.) \implies (4.) \quad \{x : f(x) \leq \alpha\} = \bigcap \{x : f(x) < \alpha + 1/n\}$$

# Measurable Functions

## Proposition (measurability condition)

Let  $f$  be an extended real-valued function on a measurable domain  $D$ . Then TFAE:

1. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) > \alpha\}$  is measurable.
2. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \geq \alpha\}$  is measurable.
3. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) < \alpha\}$  is measurable.
4. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \leq \alpha\}$  is measurable.

These imply

5. For each  $\beta \in \mathbb{R}_\infty$ , the set  $\{x : f(x) = \beta\}$  is measurable.

## Proof.

$$(1.) \implies (2.) \quad \{x : f(x) \geq \alpha\} = \bigcap \{x : f(x) > \alpha - 1/n\}$$

$$(2.) \implies (3.) \quad \{x : f(x) < \alpha\} = D - \{x : f(x) \geq \alpha\}$$

$$(3.) \implies (4.) \quad \{x : f(x) \leq \alpha\} = \bigcap \{x : f(x) < \alpha + 1/n\}$$

$$(4.) \implies (1.) \quad \{x : f(x) > \alpha\} = D - \{x : f(x) \leq \alpha\}$$

# Measurable Functions

## Proposition (measurability condition)

Let  $f$  be an extended real-valued function on a measurable domain  $D$ . Then TFAE:

1. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) > \alpha\}$  is measurable.
2. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \geq \alpha\}$  is measurable.
3. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) < \alpha\}$  is measurable.
4. For each  $\alpha \in \mathbb{R}$ , the set  $\{x : f(x) \leq \alpha\}$  is measurable.

These imply

5. For each  $\beta \in \mathbb{R}_\infty$ , the set  $\{x : f(x) = \beta\}$  is measurable.

## Proof.

$$(1.) \implies (2.) \quad \{x : f(x) \geq \alpha\} = \bigcap \{x : f(x) > \alpha - 1/n\}$$

$$(2.) \implies (3.) \quad \{x : f(x) < \alpha\} = D - \{x : f(x) \geq \alpha\}$$

$$(3.) \implies (4.) \quad \{x : f(x) \leq \alpha\} = \bigcap \{x : f(x) < \alpha + 1/n\}$$

$$(4.) \implies (1.) \quad \{x : f(x) > \alpha\} = D - \{x : f(x) \leq \alpha\}$$

$$(*) \implies (5.) \quad \text{Exercise. (2 cases: } \beta < \infty \text{ and } \beta = \pm\infty.)$$



# Definition of a Measurable Function

## Definition

Let  $D$  be measurable. Then  $f : D \rightarrow \mathbb{R}_\infty$  is measurable iff  $f$  satisfies the *measurability condition*.

## Proposition

*Let  $f$  and  $g$  be measurable (real-valued) functions defined on  $D$  and  $c \in \mathbb{R}$ . Then  $f + c$ ,  $cf$ ,  $f \pm g$ ,  $f^2$ , and  $fg$  are measurable.*



# Definition of a Measurable Function

## Definition

Let  $D$  be measurable. Then  $f : D \rightarrow \mathbb{R}_\infty$  is measurable iff  $f$  satisfies the *measurability condition*.

## Proposition

Let  $f$  and  $g$  be measurable (real-valued) functions defined on  $D$  and  $c \in \mathbb{R}$ . Then  $f + c$ ,  $cf$ ,  $f \pm g$ ,  $f^2$ , and  $fg$  are measurable.

## Proof (sketch).

$(f + c, cf)$ : Use  $\{x : f(x) + c < \alpha\} = \{x : f(x) < \alpha - c\}$ , etc.

# Definition of a Measurable Function

## Definition

Let  $D$  be measurable. Then  $f : D \rightarrow \mathbb{R}_\infty$  is measurable iff  $f$  satisfies the *measurability condition*.

## Proposition

Let  $f$  and  $g$  be measurable (real-valued) functions defined on  $D$  and  $c \in \mathbb{R}$ . Then  $f + c$ ,  $cf$ ,  $f \pm g$ ,  $f^2$ , and  $fg$  are measurable.

## Proof (sketch).

$(f + c, cf)$ : Use  $\{x : f(x) + c < \alpha\} = \{x : f(x) < \alpha - c\}$ , etc.

$(f + g)$ : If  $f(x) + g(x) < \alpha$ , there is an  $r \in \mathbb{Q}$  ( $r = r(\alpha) \neq r(x)$ ) such that  $f(x) < r < \alpha - g(x)$ . Thus

$$\{x : f(x) + g(x) < \alpha\} = \bigcup_r (\{x : f(x) < r\} \cap \{x : g(x) < \alpha - r\})$$

is a countable union of measurable sets, hence is measurable.

# Definition of a Measurable Function

## Definition

Let  $D$  be measurable. Then  $f : D \rightarrow \mathbb{R}_\infty$  is measurable iff  $f$  satisfies the *measurability condition*.

## Proposition

Let  $f$  and  $g$  be measurable (real-valued) functions defined on  $D$  and  $c \in \mathbb{R}$ . Then  $f + c$ ,  $cf$ ,  $f \pm g$ ,  $f^2$ , and  $fg$  are measurable.

## Proof (sketch).

$(f + c, cf)$ : Use  $\{x : f(x) + c < \alpha\} = \{x : f(x) < \alpha - c\}$ , etc.

$(f + g)$ : If  $f(x) + g(x) < \alpha$ , there is an  $r \in \mathbb{Q}$  ( $r = r(\alpha) \neq r(x)$ ) such that  $f(x) < r < \alpha - g(x)$ . Thus

$$\{x : f(x) + g(x) < \alpha\} = \bigcup_r (\{x : f(x) < r\} \cap \{x : g(x) < \alpha - r\})$$

is a countable union of measurable sets, hence is measurable.

$(f^2)$ : Use  $\{x : f^2(x) > \alpha\} = \{x : f(x) > \sqrt{\alpha}\} \cup \{x : f(x) < -\sqrt{\alpha}\}$ .

# Definition of a Measurable Function

## Definition

Let  $D$  be measurable. Then  $f : D \rightarrow \mathbb{R}_\infty$  is measurable iff  $f$  satisfies the *measurability condition*.

## Proposition

Let  $f$  and  $g$  be measurable (real-valued) functions defined on  $D$  and  $c \in \mathbb{R}$ . Then  $f + c$ ,  $cf$ ,  $f \pm g$ ,  $f^2$ , and  $fg$  are measurable.

## Proof (sketch).

$(f + c, cf)$ : Use  $\{x : f(x) + c < \alpha\} = \{x : f(x) < \alpha - c\}$ , etc.

$(f + g)$ : If  $f(x) + g(x) < \alpha$ , there is an  $r \in \mathbb{Q}$  ( $r = r(\alpha) \neq r(x)$ ) such that  $f(x) < r < \alpha - g(x)$ . Thus

$$\{x : f(x) + g(x) < \alpha\} = \bigcup_r (\{x : f(x) < r\} \cap \{x : g(x) < \alpha - r\})$$

is a countable union of measurable sets, hence is measurable.

$(f^2)$ : Use  $\{x : f^2(x) > \alpha\} = \{x : f(x) > \sqrt{\alpha}\} \cup \{x : f(x) < -\sqrt{\alpha}\}$ .

$(fg)$ : Use  $fg = \frac{1}{4}(f + g)^2 - \frac{1}{4}(f - g)^2$ . □

# Sequences of Measurable Functions

## Theorem

Let  $\{f_n\}$  be a sequence of measurable functions on a common domain  $D$ . Then the functions

$$\sup\{f_1, \dots, f_n\}, \quad \sup_n f_n, \quad \limsup_n f_n$$

are measurable. Analogous statements hold for  $\inf$  and  $\liminf$ .

# Sequences of Measurable Functions

## Theorem

Let  $\{f_n\}$  be a sequence of measurable functions on a common domain  $D$ . Then the functions

$$\sup\{f_1, \dots, f_n\}, \quad \sup_n f_n, \quad \limsup_n f_n$$

are measurable. Analogous statements hold for  $\inf$  and  $\liminf$ .

## Proof.

Set  $h = \sup\{f_1, \dots, f_n\}$ , then

$$\{x : h(x) > \alpha\} = \bigcup_{i=1}^n \{x : f_i(x) > \alpha\}.$$

Hence  $h$  is measurable. Now set  $g = \sup_n f_n$ , then

$$\{x : g(x) > \alpha\} = \bigcup_{i=1}^{\infty} \{x : f_i(x) > \alpha\}.$$

Hence  $g$  is measurable. Combine the above with the definition

$\limsup_n f_n = \inf_n \sup_{k \geq n} f_k$  to finish. □

# 'Simple' Functions are Measurable

## Proposition

If  $f$  is measurable and  $g = f$  a.e., then  $g$  is measurable.

## Proof.

Set  $E = \{x : f(x) \neq g(x)\}$ . Then  $m(E) = 0$ . So  $\{x : g(x) > \alpha\} = \{x : f(x) > \alpha\} \cup \{x \in E : g(x) > \alpha\} - \{x \in E : g(x) \leq \alpha\}$ .  $\square$

## Definition

A measurable real-valued function  $\phi$  is *simple* if it assumes only finitely many values. Then

$$\phi(x) = \sum_{k=1}^n \alpha_k \chi_{A_k}(x) \quad \text{where} \quad A_k = \{x : \phi(x) = \alpha_k\}$$

If each  $A_k$  is an interval, then  $\phi$  is called a *step function*.

# 'Simple' Functions are Measurable

## Proposition

If  $f$  is measurable and  $g = f$  a.e., then  $g$  is measurable.

## Proof.

Set  $E = \{x : f(x) \neq g(x)\}$ . Then  $m(E) = 0$ . So  $\{x : g(x) > \alpha\} = \{x : f(x) > \alpha\} \cup \{x \in E : g(x) > \alpha\} - \{x \in E : g(x) \leq \alpha\}$ .  $\square$

## Definition

A measurable real-valued function  $\phi$  is *simple* if it assumes only finitely many values. Then

$$\phi(x) = \sum_{k=1}^n \alpha_k \chi_{A_k}(x) \quad \text{where} \quad A_k = \{x : \phi(x) = \alpha_k\}$$

If each  $A_k$  is an interval, then  $\phi$  is called a *step function*.

## Example

►  $s(x) = \sum_{k=1}^N \frac{k^2}{N^2} \chi_{[\frac{k-1}{N}, \frac{k}{N}]}(x)$  is a step function;  $\chi_{\mathbb{Q}}$  is simple.



# Measurable Functions are 'Simple'

## Proposition

*Let  $f : [a, b] \rightarrow \mathbb{R}_\infty$  be measurable such that  $m(\{f(x) = \pm\infty\})$  is zero. Given  $\epsilon > 0$ , there is a step function  $s$  and a continuous function  $h$  so that  $|f - s| < \epsilon$  and  $|f - h| < \epsilon$  a.e.*

# Measurable Functions are 'Simple'

## Proposition

Let  $f: [a, b] \rightarrow \mathbb{R}_\infty$  be measurable such that  $m(\{f(x) = \pm\infty\})$  is zero. Given  $\epsilon > 0$ , there is a step function  $s$  and a continuous function  $h$  so that  $|f - s| < \epsilon$  and  $|f - h| < \epsilon$  a.e.

## Proof (Exercise).

1. There is an  $M$  such that  $|f| \leq M$  except on a set of measure  $< \epsilon/3$ .
2. There is a simple function  $\phi$  such that  $|f - \phi| < \epsilon$  except when  $|f| > M$ . (Hint:  $(M - -M) \leq n \cdot \epsilon$ .)
3. There is a step function  $g$  such that  $g = \phi$  except on a set of measure  $< \epsilon/3$ . (Hint: look [here](#).)
4. There is a continuous function  $h$  such that  $h = g$  except on a set of measure  $< \epsilon/3$ . (Hint: think like a spline.)



# Functionally Measured Exercises

## Exercises

1. Let  $\phi_1$  and  $\phi_2$  be simple functions and  $c \in \mathbb{R}$ . Show that
  - a.  $c\phi$  is a simple function,
  - b.  $\phi_1 + \phi_2$  is a simple function,
  - c.  $\phi_1 \cdot \phi_2$  is a simple function.
2. For a set  $S$  define the characteristic or indicator function to be  $\chi_S(x) = \begin{cases} 1 & x \in S \\ 0 & x \notin S \end{cases}$ . Show that
  - a.  $\chi_{A \cap B} = \chi_A \cdot \chi_B$ ,
  - b.  $\chi_{A \cup B} = \chi_A + \chi_B - \chi_A \cdot \chi_B$ .
  - c.  $\chi_{A^c} = 1 - \chi_A$ .
3. Let  $D$  be a dense set of real numbers; i.e., every interval contains an element of  $D$ . Let  $f$  be an extended real-valued function on  $\mathbb{R}$  such that for any  $d \in D$ , the set  $\{x : f(x) > d\}$  is measurable. Then  $f$  is measurable.

# Integration

We began by looking at two examples of integration problems.

- ▶ The Riemann integral over  $[0, 1]$  of a function with infinitely many discontinuities didn't exist even though the points of discontinuity formed a set of measure zero. (The points of discontinuity formed a dense set in  $[0, 1]$ .)
- ▶ The limit of a sequence of Riemann integrable functions did not equal the integral of the limit function of the sequence. (Each function had area  $1/2$ , but the limit of the sequence was the zero function.)

We will look at Riemann integration, then Riemann-Stieltjes integration, and last, develop the Lebesgue integral.

There are many other types of integrals: Darboux, Denjoy, Gauge, Perron, etc. See the list given in the “See also” section of *Integrals* on Mathworld.

# Riemann Integral

## Definition

- ▶ A *partition*  $\mathcal{P}$  of  $[a, b]$  is a finite set of points such that  $\mathcal{P} = \{a = x_0 < x_1 < \cdots < x_{n-1} < x_n = b\}$ .
- ▶ Set  $M_i = \sup f(x)$  on  $[x_{i-1}, x_i]$ . The *upper sum* of  $f$  on  $[a, b]$  w.r.t.  $\mathcal{P}$  is

$$U(\mathcal{P}, f) = \sum_{i=1}^n M_i \cdot \Delta x_i$$

- ▶ The *upper Riemann integral* of  $f$  over  $[a, b]$  is

$$\int_a^b f(x) dx = \inf_{\mathcal{P}} U(\mathcal{P}, f)$$

## Exercise

1. Define the lower sum  $L(\mathcal{P}, f)$  and the lower integral  $\int_a^b f$ .

# Definitely a Riemann Integral

## Definition

If  $\int_a^b f(x) dx = \int_a^b f(x) dx$ , then  $f$  is Riemann integrable and is written as  $\int_a^b f(x) dx$  and  $f \in \mathfrak{R}$  on  $[a, b]$ .

## Proposition

*A function  $f$  is Riemann integrable on  $[a, b]$  if and only if for every  $\epsilon > 0$  there is a partition  $\mathcal{P}$  of  $[a, b]$  such that*

$$U(\mathcal{P}, f) - L(\mathcal{P}, f) < \epsilon.$$

# Definitely a Riemann Integral

## Definition

If  $\int_a^b f(x) dx = \int_a^b f(x) dx$ , then  $f$  is Riemann integrable and is written as  $\int_a^b f(x) dx$  and  $f \in \mathfrak{R}$  on  $[a, b]$ .

## Proposition

*A function  $f$  is Riemann integrable on  $[a, b]$  if and only if for every  $\epsilon > 0$  there is a partition  $\mathcal{P}$  of  $[a, b]$  such that*

$$U(\mathcal{P}, f) - L(\mathcal{P}, f) < \epsilon.$$

## Theorem

*If  $f$  is continuous on  $[a, b]$ , then  $f \in \mathfrak{R}$  on  $[a, b]$ .*

# Definitely a Riemann Integral

## Definition

If  $\int_a^b f(x) dx = \int_a^b f(x) dx$ , then  $f$  is Riemann integrable and is written as  $\int_a^b f(x) dx$  and  $f \in \mathfrak{R}$  on  $[a, b]$ .

## Proposition

*A function  $f$  is Riemann integrable on  $[a, b]$  if and only if for every  $\epsilon > 0$  there is a partition  $\mathcal{P}$  of  $[a, b]$  such that*

$$U(\mathcal{P}, f) - L(\mathcal{P}, f) < \epsilon.$$

## Theorem

*If  $f$  is continuous on  $[a, b]$ , then  $f \in \mathfrak{R}$  on  $[a, b]$ .*

## Theorem

*If  $f$  is bounded on  $[a, b]$  with only finitely many points of discontinuity, then  $f \in \mathfrak{R}$  on  $[a, b]$ .*



# Properties of Riemann Integrals

## Proposition

Let  $f$  and  $g \in \mathfrak{R}$  on  $[a, b]$  and  $c \in \mathbb{R}$ . Then

- ▶  $\int_a^b cf \, dx = c \int_a^b f \, dx$
- ▶  $\int_a^b (f + g) \, dx = \int_a^b f \, dx + \int_a^b g \, dx$
- ▶  $f \cdot g \in \mathfrak{R}$
- ▶ if  $f \leq g$ , then  $\int_a^b f \, dx \leq \int_a^b g \, dx$

# Properties of Riemann Integrals

## Proposition

Let  $f$  and  $g \in \mathfrak{R}$  on  $[a, b]$  and  $c \in \mathbb{R}$ . Then

- ▶  $\int_a^b cf \, dx = c \int_a^b f \, dx$
- ▶  $\int_a^b (f + g) \, dx = \int_a^b f \, dx + \int_a^b g \, dx$
- ▶  $f \cdot g \in \mathfrak{R}$
- ▶ if  $f \leq g$ , then  $\int_a^b f \, dx \leq \int_a^b g \, dx$
  
- ▶  $\left| \int_a^b f \, dx \right| \leq \int_a^b |f| \, dx$

# Properties of Riemann Integrals

## Proposition

Let  $f$  and  $g \in \mathfrak{R}$  on  $[a, b]$  and  $c \in \mathbb{R}$ . Then

- ▶  $\int_a^b cf \, dx = c \int_a^b f \, dx$
- ▶  $\int_a^b (f + g) \, dx = \int_a^b f \, dx + \int_a^b g \, dx$
- ▶  $f \cdot g \in \mathfrak{R}$
- ▶ if  $f \leq g$ , then  $\int_a^b f \, dx \leq \int_a^b g \, dx$
  
- ▶  $\left| \int_a^b f \, dx \right| \leq \int_a^b |f| \, dx$
  
- ▶ Define  $F(x) = \int_a^x f(t) \, dt$ . Then  $F$  is continuous and, if  $f$  is continuous at  $x_0$ , then  $F'(x_0) = f(x_0)$

# Properties of Riemann Integrals

## Proposition

Let  $f$  and  $g \in \mathfrak{R}$  on  $[a, b]$  and  $c \in \mathbb{R}$ . Then

- ▶  $\int_a^b cf \, dx = c \int_a^b f \, dx$
- ▶  $\int_a^b (f + g) \, dx = \int_a^b f \, dx + \int_a^b g \, dx$
- ▶  $f \cdot g \in \mathfrak{R}$
- ▶ if  $f \leq g$ , then  $\int_a^b f \, dx \leq \int_a^b g \, dx$
  
- ▶  $\left| \int_a^b f \, dx \right| \leq \int_a^b |f| \, dx$
  
- ▶ Define  $F(x) = \int_a^x f(t) \, dt$ . Then  $F$  is continuous and, if  $f$  is continuous at  $x_0$ , then  $F'(x_0) = f(x_0)$
  
- ▶ If  $F' = f$  on  $[a, b]$ , then  $\int_a^b f(x) \, dx = F(b) - F(a)$

# Riemann Integrated Exercises

## Exercises

1. If  $\int_a^b |f(x)| dx = 0$ , then  $f = 0$ .
2. Show why  $\int_0^1 \chi_{\mathbb{Q}}(x) dx$  does not exist.
3. Define

$$S_n(x) = \sum_{k=1}^{n+1} \left( \frac{k-1}{k} \cdot \chi_{\left[\frac{k-1}{k}, \frac{k}{k+1}\right)}(x) \right) + \frac{n}{n+1} \chi_{\left[\frac{n+1}{n+2}, 1\right]}(x).$$

- 3.1 How many discontinuities does  $S_n$  have?
- 3.2 Prove that  $S'_n(x) = 0$  a.e.
- 3.3 Calculate  $\int_0^1 S_n(x) dx$ .
- 3.4 What is  $S_\infty$ ?
- 3.5 Does  $\int_0^1 S_\infty(x) dx$  exist?

(See an animated graph of  $S_N$ .)

# Riemann-Stieltjes Integral

## Definition

- ▶ Let  $\alpha(x)$  be a monotonically increasing function on  $[a, b]$ . Set  $\Delta\alpha_i = \alpha(x_i) - \alpha(x_{i-1})$ .
- ▶ Set  $M_i = \sup f(x)$  on  $[x_{i-1}, x_i]$ . The *upper sum* of  $f$  on  $[a, b]$  w.r.t.  $\alpha$  and  $\mathcal{P}$  is

$$U(\mathcal{P}, f, \alpha) = \sum_{i=1}^n M_i \cdot \Delta\alpha_i$$

- ▶ The *upper Riemann-Stieltjes integral* of  $f$  over  $[a, b]$  w.r.t.  $\alpha$  is

$$\int_a^b f(x) d\alpha(x) = \inf_{\mathcal{P}} U(\mathcal{P}, f, \alpha)$$

## Exercise

1. Define the lower sum  $L(\mathcal{P}, f, \alpha)$  and lower integral  $\int_a^b f d\alpha$ .

# Definitely a Riemann-Stieltjes Integral

## Definition

If  $\int_a^b f d\alpha = \int_a^b f d\alpha$ , then  $f$  is Riemann-Stieltjes integrable and is written as  $\int_a^b f(x) d\alpha(x)$  and  $f \in \mathfrak{R}(\alpha)$  on  $[a, b]$ .

## Proposition

*A function  $f$  is Riemann-Stieltjes integrable w.r.t.  $\alpha$  on  $[a, b]$  iff for every  $\epsilon > 0$  there is a partition  $\mathcal{P}$  of  $[a, b]$  such that*

$$U(\mathcal{P}, f, \alpha) - L(\mathcal{P}, f, \alpha) < \epsilon.$$

# Definitely a Riemann-Stieltjes Integral

## Definition

If  $\int_a^b f d\alpha = \int_a^b f d\alpha$ , then  $f$  is Riemann-Stieltjes integrable and is written as  $\int_a^b f(x) d\alpha(x)$  and  $f \in \mathfrak{R}(\alpha)$  on  $[a, b]$ .

## Proposition

*A function  $f$  is Riemann-Stieltjes integrable w.r.t.  $\alpha$  on  $[a, b]$  iff for every  $\epsilon > 0$  there is a partition  $\mathcal{P}$  of  $[a, b]$  such that*

$$U(\mathcal{P}, f, \alpha) - L(\mathcal{P}, f, \alpha) < \epsilon.$$

## Theorem

*If  $f$  is continuous on  $[a, b]$ , then  $f \in \mathfrak{R}(\alpha)$  on  $[a, b]$ .*



# Definitely a Riemann-Stieltjes Integral

## Definition

If  $\int_a^b f d\alpha = \int_a^b f d\alpha$ , then  $f$  is Riemann-Stieltjes integrable and is written as  $\int_a^b f(x) d\alpha(x)$  and  $f \in \mathfrak{R}(\alpha)$  on  $[a, b]$ .

## Proposition

*A function  $f$  is Riemann-Stieltjes integrable w.r.t.  $\alpha$  on  $[a, b]$  iff for every  $\epsilon > 0$  there is a partition  $\mathcal{P}$  of  $[a, b]$  such that*

$$U(\mathcal{P}, f, \alpha) - L(\mathcal{P}, f, \alpha) < \epsilon.$$

## Theorem

*If  $f$  is continuous on  $[a, b]$ , then  $f \in \mathfrak{R}(\alpha)$  on  $[a, b]$ .*

## Theorem

*If  $f$  is bounded on  $[a, b]$  with only finitely many points of discontinuity and  $\alpha$  is continuous at each of  $f$ 's discontinuities, then  $f \in \mathfrak{R}(\alpha)$  on  $[a, b]$ .*

# Properties of Riemann-Stieltjes Integrals

## Proposition

Let  $f$  and  $g \in \mathfrak{R}(\alpha)$  and in  $\beta$  on  $[a, b]$  and  $c \in \mathbb{R}$ . Then

- ▶  $\int_a^b c f d\alpha = c \int_a^b f d\alpha$  and  $\int_a^b f d(c\alpha) = c \int_a^b f d\alpha$
- ▶  $\int_a^b (f + g) d\alpha = \int_a^b f d\alpha + \int_a^b g d\alpha$  and  
 $\int_a^b f d(\alpha + \beta) = \int_a^b f d\alpha + \int_a^b f d\beta$
- ▶  $f \cdot g \in \mathfrak{R}(\alpha)$
- ▶ if  $f \leq g$ , then  $\int_a^b f d\alpha \leq \int_a^b g d\alpha$

# Properties of Riemann-Stieltjes Integrals

## Proposition

Let  $f$  and  $g \in \mathfrak{R}(\alpha)$  and in  $\beta$  on  $[a, b]$  and  $c \in \mathbb{R}$ . Then

- ▶  $\int_a^b c f d\alpha = c \int_a^b f d\alpha$  and  $\int_a^b f d(c\alpha) = c \int_a^b f d\alpha$
- ▶  $\int_a^b (f + g) d\alpha = \int_a^b f d\alpha + \int_a^b g d\alpha$  and  
 $\int_a^b f d(\alpha + \beta) = \int_a^b f d\alpha + \int_a^b f d\beta$
- ▶  $f \cdot g \in \mathfrak{R}(\alpha)$
- ▶ if  $f \leq g$ , then  $\int_a^b f d\alpha \leq \int_a^b g d\alpha$
- ▶  $\left| \int_a^b f d\alpha \right| \leq \int_a^b |f| d\alpha$

# Properties of Riemann-Stieltjes Integrals

## Proposition

Let  $f$  and  $g \in \mathfrak{R}(\alpha)$  and in  $\beta$  on  $[a, b]$  and  $c \in \mathbb{R}$ . Then

▶  $\int_a^b c f d\alpha = c \int_a^b f d\alpha$  and  $\int_a^b f d(c\alpha) = c \int_a^b f d\alpha$

▶  $\int_a^b (f + g) d\alpha = \int_a^b f d\alpha + \int_a^b g d\alpha$  and

$\int_a^b f d(\alpha + \beta) = \int_a^b f d\alpha + \int_a^b f d\beta$

▶  $f \cdot g \in \mathfrak{R}(\alpha)$

▶ if  $f \leq g$ , then  $\int_a^b f d\alpha \leq \int_a^b g d\alpha$

▶  $\left| \int_a^b f d\alpha \right| \leq \int_a^b |f| d\alpha$

▶ Suppose that  $\alpha' \in \mathfrak{R}$  and  $f$  is bounded. Then  $f \in \mathfrak{R}(\alpha)$  iff  $f\alpha' \in \mathfrak{R}$  and

$$\int_a^b f d\alpha = \int_a^b f \cdot \alpha' dx$$

# Riemann-Stieltjes Integrals and Series

## Proposition

If  $f$  is continuous at  $c \in (a, b)$  and  $\alpha(x) = r$  for  $a \leq x < c$  and  $\alpha(x) = s$  for  $c < x \leq b$ , then

$$\begin{aligned}\int_a^b f d\alpha &= f(c) (\alpha(c+) - \alpha(c-)) \\ &= f(c) (s - r)\end{aligned}$$

## Proposition

Let  $\alpha = [x]$ , the greatest integer function. If  $f$  is continuous on  $[0, b]$ , then

$$\int_0^b f(x) d[x] = \sum_{k=1}^{\lfloor b \rfloor} f(k)$$

# Riemann-Stieltjes Integrated Exercises

## Exercises

1.  $\int_0^1 x dx^2$

2.  $\int_0^{\pi/2} \cos(x) d \sin(x)$

3.  $\int_0^{5/2} x d(x - \lfloor x \rfloor)$

4.  $\int_{-1}^1 e^x d|x|$

5.  $\int_{-3/2}^{3/2} e^x d\lfloor x \rfloor$

6.  $\int_{-1}^1 e^x d\lfloor x \rfloor$

7. Set  $H$  to be the Heaviside function; i.e.,

$$H(x) = \begin{cases} 0 & x \leq 0 \\ 1 & \text{otherwise} \end{cases}.$$

Show that, if  $f$  is continuous at 0, then

$$\int_{-\infty}^{+\infty} f(x) dH(x) = f(0).$$

# Lebesgue Integral

We start with **simple functions**.

## Definition

A function has *finite support* if it vanishes outside a finite interval.

## Definition

Let  $\phi$  be a measurable simple function with finite support. If

$\phi(x) = \sum_{i=1}^n a_i \chi_{A_i}(x)$  is a representation of  $\phi$ , then

$$\int \phi(x) dx = \sum_{i=1}^n a_i \cdot m(A_i)$$

## Definition

If  $E$  is a measurable set, then  $\int_E \phi = \int \phi \cdot \chi_E$ .

# Integral Linearity

## Proposition

If  $\phi$  and  $\psi$  are measurable simple functions with finite support and  $a, b \in \mathbb{R}$ , then  $\int (a\phi + b\psi) = a \int \phi + b \int \psi$ . Further, if  $\phi \leq \psi$  a.e., then  $\int \phi \leq \int \psi$ .

## Proof (sketch).

- I. Let  $\phi = \sum^N \alpha_i \chi_{A_i}$  and  $\psi = \sum^M \beta_i \chi_{B_i}$ . Then show  $a\phi + b\psi$  can be written as  $a\phi + b\psi = \sum^K (a\alpha_{k_i} + b\beta_{k_j}) \chi_{E_k}$  for the properly chosen  $E_k$ . Set  $A_0$  and  $B_0$  to be zero sets of  $\phi$  and  $\psi$ . (Take  $\{E_k : k = 0..K\} = \{A_j \cap B_k : j = 0..N, k = 0..M\}$ .)
- II. Use the definition to show  $\int \psi - \int \phi = \int (\psi - \phi) \geq \int 0 = 0$ .  $\square$



# Steps to the Lebesgue Integral

## Proposition

Let  $f$  be bounded on  $E \in \mathfrak{M}$  with  $m(E) < \infty$ . Then  $f$  is measurable iff

$$\inf_{f \leq \psi} \int_E \psi = \sup_{f \geq \phi} \int_E \phi$$

for all simple functions  $\phi$  and  $\psi$ .

# Steps to the Lebesgue Integral

## Proposition

Let  $f$  be bounded on  $E \in \mathfrak{M}$  with  $m(E) < \infty$ . Then  $f$  is measurable iff

$$\inf_{f \leq \psi} \int_E \psi = \sup_{f \geq \phi} \int_E \phi$$

for all simple functions  $\phi$  and  $\psi$ .

## Proof.

I. Suppose  $f$  is bounded by  $m$ . Define

$$E_k = \left\{ x : \frac{k-1}{n}M < f(x) \leq \frac{k}{n}M \right\}, \quad -n \leq k \leq n$$

The  $E_k$  are measurable, disjoint, and have union  $E$ . Set

$$\psi_n(x) = \frac{M}{n} \sum_{-n}^n k \chi_{E_k}(x), \quad \phi_n(x) = \frac{M}{n} \sum_{-n}^n (k-1) \chi_{E_k}(x)$$

## SLI (cont)

(proof cont).

Then  $\phi_n(x) \leq f(x) \leq \psi(x)$ , and so

$$\blacktriangleright \inf \int_E \psi \leq \int_E \psi_n = \frac{M}{n} \sum_{k=-n}^n k m(E_k)$$


$$\blacktriangleright \sup \int_E \phi \geq \int_E \phi_n = \frac{M}{n} \sum_{k=-n}^n (k-1) m(E_k)$$

Thus  $0 \leq \inf \int_E \psi - \sup \int_E \phi \leq \frac{M}{n} m(E)$ . Since  $n$  is arbitrary, equality holds.

II. Suppose that  $\inf \int_E \psi = \sup \int_E \phi$ . Choose  $\phi_n$  and  $\psi_n$  so that  $\phi_n \leq f \leq \psi_n$  and  $\int_E (\psi_n - \phi_n) < \frac{1}{n}$ . The functions  $\psi^* = \inf \psi_n$  and  $\phi^* = \sup \phi_n$  are measurable and  $\phi^* \leq f \leq \psi^*$ . The set  $\Delta = \{x : \phi^*(x) < \psi^*(x)\}$  has measure 0. Thus  $\phi^* = \psi^*$  almost everywhere, so  $\phi^* = f$  a.e. Hence  $f$  is measurable.  $\square$

# Example Steps

## Example



# Defining the Lebesgue Integral

## Definition

If  $f$  is a bounded measurable function on a measurable set  $E$  with  $m(E) < \infty$ , then

$$\int_E f = \inf_{\psi \geq f} \int_E \psi$$

for all simple functions  $\psi \geq f$ .

# Defining the Lebesgue Integral

## Definition

If  $f$  is a bounded measurable function on a measurable set  $E$  with  $m(E) < \infty$ , then

$$\int_E f = \inf_{\psi \geq f} \int_E \psi$$

for all simple functions  $\psi \geq f$ .

## Proposition

*Let  $f$  be a bounded function defined on  $E = [a, b]$ . If  $f$  is Riemann integrable on  $[a, b]$ , then  $f$  is measurable on  $[a, b]$  and*

$$\int_E f = \int_a^b f(x) dx;$$

*the Riemann integral of  $f$  equals the Lebesgue integral of  $f$ .*

# Properties of the Lebesgue Integral

## Proposition

If  $f$  and  $g$  are measurable on  $E$ , a set of finite measure, then

$$\blacktriangleright \int_E (\alpha f + \beta g) = \alpha \int_E f + \beta \int_E g$$

$$\blacktriangleright \text{if } f = g \text{ a.e., then } \int_E f = \int_E g$$

$$\blacktriangleright \text{if } f \leq g \text{ a.e., then } \int_E f \leq \int_E g$$

$$\blacktriangleright \left| \int_E f \right| \leq \int_E |f|$$

$$\blacktriangleright \text{if } a \leq f \leq b, \text{ then } a \cdot m(E) \leq \int_E f \leq b \cdot m(E)$$

$$\blacktriangleright \text{if } A \cap B = \emptyset, \text{ then } \int_{A \cup B} f = \int_A f + \int_B f$$

Proof.

Exercise.



# Lebesgue Integral Examples

## Examples

1. Let  $D(x) = \begin{cases} \frac{1}{q} & x = \frac{p}{q} \in \mathbb{Q} \\ 0 & \text{otherwise} \end{cases}$ . Then  $\int_{[0,1]} D = \int_0^1 D(x) dx$ .

2. Let  $\chi_{\mathbb{Q}}(x) = \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & \text{otherwise} \end{cases}$ . Then  $\int_{[0,1]} \chi_{\mathbb{Q}} \neq \int_0^1 \chi_{\mathbb{Q}}(x) dx$ .

3. Define

$$f_n(x) = \sum_{k=1}^{n+1} \left( \frac{k-1}{k} \cdot \chi_{\left[\frac{k-1}{k}, \frac{k}{k+1}\right)}(x) \right) + \frac{n}{n+1} \chi_{\left[\frac{n+1}{n+2}, 1\right]}(x).$$

Then

3.1  $f_n$  is a step function, hence integrable

3.2  $f'_n(x) = 0$  a.e.

3.3  $\frac{1}{4} \leq \int_{[0,1]} f_n = \int_0^1 f_n(x) dx < \frac{3}{8}$



# Extending the Integral Definition

## Definition

Let  $f$  be a nonnegative measurable function defined on a measurable set  $E$ . Define

$$\int_E f = \sup_{h \leq f} \int_E h$$

where  $h$  is a bounded measurable function with finite support.

## Proposition

*If  $f$  and  $g$  are nonnegative measurable functions, then*

- ▶  $\int_E c f = c \int_E f$  for  $c > 0$
- ▶  $\int_E f + g = \int_E f + \int_E g$
- ▶ *If  $f \leq g$  a.e., then  $\int_E f \leq \int_E g$*

## Proof.

Exercise.

# General Lebesgue's Integral

## Definition

Set  $f^+(x) = \max\{f(x), 0\}$  and  $f^-(x) = \max\{-f(x), 0\}$ . Then  $f = f^+ - f^-$  and  $|f| = f^+ + f^-$ . A measurable function  $f$  is integrable over  $E$  iff both  $f^+$  and  $f^-$  are integrable over  $E$ , and then  $\int_E f = \int_E f^+ - \int_E f^-$ .

## Proposition

Let  $f$  and  $g$  be integrable over  $E$  and let  $c \in \mathbb{R}$ . Then

1.  $\int_E cf = c \int_E f$
2.  $\int_E f + g = \int_E f + \int_E g$
3. if  $f \leq g$  a.e., then  $\int_E f \leq \int_E g$
4. if  $A, B$  are disjoint m'ble subsets of  $E$ ,  $\int_{A \cup B} f = \int_A f + \int_B f$

# Convergence Theorems

## Theorem (Bounded Convergence Theorem)

Let  $\{f_n : E \rightarrow \mathbb{R}\}$  be a sequence of measurable functions converging to  $f$  with  $m(E) < \infty$ . If there is a uniform bound  $M$  for all  $f_n$ , then

$$\int_E \lim_n f_n = \lim_n \int_E f_n$$

# Convergence Theorems

## Theorem (Bounded Convergence Theorem)

Let  $\{f_n : E \rightarrow \mathbb{R}\}$  be a sequence of measurable functions converging to  $f$  with  $m(E) < \infty$ . If there is a uniform bound  $M$  for all  $f_n$ , then

$$\int_E \lim_n f_n = \lim_n \int_E f_n$$

### Proof (sketch).

Let  $\epsilon > 0$ .

1.  $f_n$  converges “almost uniformly;” i.e.,  $\exists A, N$  s.t.  $m(A) < \frac{\epsilon}{4M}$   
and, for  $n > N, x \in E - A \implies |f_n(x) - f(x)| \leq \frac{\epsilon}{2m(E)}$ .

2.  $\left| \int_E f_n - \int_E f \right| = \left| \int_E f_n - f \right| \leq \int_E |f_n - f| = \left( \int_{E-A} + \int_A \right) |f_n - f|$

3.  $\int_{E-A} |f_n - f| + \int_A |f_n| + |f| \leq \frac{\epsilon}{2m(E)} \cdot m(E) + 2M \cdot \frac{\epsilon}{4M} = \epsilon \quad \square$

# Lebesgue's Dominated Convergence Theorem

## Theorem (Dominated Convergence Theorem)

*Let  $\{f_n : E \rightarrow \mathbb{R}\}$  be a sequence of measurable functions converging a.e. on  $E$  with  $m(E) < \infty$ . If there is an integrable function  $g$  on  $E$  such that  $|f_n| \leq g$  then*

$$\int_E \lim_n f_n = \lim_n \int_E f_n$$

# Lebesgue's Dominated Convergence Theorem

## Theorem (Dominated Convergence Theorem)

Let  $\{f_n : E \rightarrow \mathbb{R}\}$  be a sequence of measurable functions converging a.e. on  $E$  with  $m(E) < \infty$ . If there is an integrable function  $g$  on  $E$  such that  $|f_n| \leq g$  then

$$\int_E \lim_n f_n = \lim_n \int_E f_n$$

## Lemma

Under the conditions of the DCT, set  $g_n = \sup_{k \geq n} \{f_n, f_{n+1}, \dots\}$

and  $h_n = \inf_{k \geq n} \{f_n, f_{n+1}, \dots\}$ . Then  $g_n$  and  $h_n$  are integrable and

$\lim g_n = f = \lim h_n$  a.e.

# Lebesgue's Dominated Convergence Theorem

## Theorem (Dominated Convergence Theorem)

Let  $\{f_n : E \rightarrow \mathbb{R}\}$  be a sequence of measurable functions converging a.e. on  $E$  with  $m(E) < \infty$ . If there is an integrable function  $g$  on  $E$  such that  $|f_n| \leq g$  then

$$\int_E \lim_n f_n = \lim_n \int_E f_n$$

## Lemma

Under the conditions of the DCT, set  $g_n = \sup_{k \geq n} \{f_n, f_{n+1}, \dots\}$

and  $h_n = \inf_{k \geq n} \{f_n, f_{n+1}, \dots\}$ . Then  $g_n$  and  $h_n$  are integrable and  $\lim g_n = f = \lim h_n$  a.e.

## Proof of DCT (sketch).

- ▶ Both  $g_n$  and  $h_n$  are monotone and converging. Apply MCT.
- ▶  $h_n \leq f_n \leq g_n \implies \int_E h_n \leq \int_E f_n \leq \int_E g_n$ . □

# Increasing the Convergence

## Theorem (Fatou's Lemma)

If  $\{f_n\}$  is a sequence of measurable functions converging to  $f$  a.e. on  $E$ , then

$$\int_E \liminf_n f_n \leq \liminf_n \int_E f_n$$



# Increasing the Convergence

## Theorem (Fatou's Lemma)

If  $\{f_n\}$  is a sequence of measurable functions converging to  $f$  a.e. on  $E$ , then

$$\int_E \liminf_n f_n \leq \liminf_n \int_E f_n$$

## Theorem (Monotone Convergence Theorem)

If  $\{f_n\}$  is an increasing sequence of nonnegative measurable functions converging to  $f$ , then

$$\int \lim_n f_n = \lim_n \int f_n$$

# Increasing the Convergence

## Theorem (Fatou's Lemma)

If  $\{f_n\}$  is a sequence of measurable functions converging to  $f$  a.e. on  $E$ , then

$$\int_E \liminf_n f_n \leq \liminf_n \int_E f_n$$

## Theorem (Monotone Convergence Theorem)

If  $\{f_n\}$  is an increasing sequence of nonnegative measurable functions converging to  $f$ , then

$$\int \lim_n f_n = \lim_n \int f_n$$

## Corollary (Beppo Levi Theorem (cf.))

If  $\{f_n\}$  is a sequence of nonnegative measurable functions, then

$$\int \sum_{n=1}^{\infty} f_n = \sum_{n=1}^{\infty} \int f_n$$

## Sidebar: Littlewood's Three Principles

John Edensor Littlewood said,

*The extent of knowledge required is nothing so great as sometimes supposed. There are three principles, roughly expressible in the following terms:*

- ▶ *every measurable set is nearly a finite union of intervals;*
- ▶ *every measurable function is nearly continuous;*
- ▶ *every convergent sequence of measurable functions is nearly uniformly convergent.*

*Most of the results of analysis are fairly intuitive applications of these ideas.*

From *Lectures on the Theory of Functions*, Oxford, 1944, p. 26.

# Extensions of Convergence

The sequence  $f_n$  converges to  $f \dots$

**Definition (Convergence Almost Everywhere)**

*almost everywhere* if  $m(\{x : f_n(x) \not\rightarrow f(x)\}) = 0$ .

**Definition (Convergence Almost Uniformly)**

*almost uniformly on  $E$*  if, for any  $\epsilon > 0$ , there is a set  $A \subset E$  with  $m(A) < \epsilon$  so that  $f_n$  converges uniformly on  $E - A$ .

**Definition (Convergence in Measure)**

*in measure* if, for any  $\epsilon > 0$ ,  $\lim_{n \rightarrow \infty} m(\{x : |f_n(x) - f(x)| \geq \epsilon\}) = 0$ .

**Definition (Convergence in Mean (of order  $p > 1$ ))**

*in mean* if  $\lim_{n \rightarrow \infty} \|f_n - f\|_p = \lim_{n \rightarrow \infty} \left[ \int_E |f - f_n|^p \right]^{1/p} = 0$

# Integrated Exercises

## Exercises

1. Prove: If  $f$  is integrable on  $E$ , then  $|f|$  is integrable on  $E$ .
2. Prove: If  $f$  is integrable over  $E$ , then  $\left| \int_E f \right| \leq \int_E |f|$ .
3. True or False: If  $|f|$  is integrable over  $E$ , then  $f$  is integrable over  $E$ .
4. Let  $f$  be integrable over  $E$ . For any  $\epsilon > 0$ , there is a simple (resp. step) function  $\phi$  (resp.  $\psi$ ) such that  $\int_E |f - \phi| < \epsilon$ .
5. For  $n = k + 2^\nu$ ,  $0 \leq k < 2^\nu$ , define  $f_n = \chi_{[k2^{-\nu}, (k+1)2^{-\nu}]}$ .
  - 5.1 Show that  $f_n$  does not converge for any  $x \in [0, 1]$ .
  - 5.2 Show that  $f_n$  does not converge a.e. on  $[0, 1]$ .
  - 5.3 Show that  $f_n$  does not converge almost uniformly on  $[0, 1]$ .
  - 5.4 Show that  $f_n \rightarrow 0$  in measure.
  - 5.5 Show that  $f_n \rightarrow 0$  in mean (of order 2).

# References

Texts on analysis, integration, and measure:

- ▶ *Mathematical Analysis*, T. Apostle
- ▶ *Principles of Mathematical Analysis*, W. Rudin
- ▶ *Real Analysis*, H. Royden
- ▶ *Lebesgue Integration*, S. Chae
- ▶ *Geometric Measure Theory*, F. Morgan

Comparison of different types of integrals:

- ▶ *Integral, Measure, and Derivative: A Unified Approach*, G. Shilov and B. Gurevich